

- 1 -

## FLAVONOID 3',5' HYDROXYLASE GENE SEQUENCES AND USES THEREFOR

### BACKGROUND OF THE INVENTION

#### 5 FIELD OF THE INVENTION

The present invention relates generally to a genetic sequence encoding a polypeptide having flavonoid 3', 5'-hydroxylase (F3'5'H) activity and to the use of the genetic sequence and/or its corresponding polypeptide thereof *inter alia* to manipulate color in flowers or  
10 parts thereof or in other plant tissue. More particularly, the F3'5'H has the ability to modulate dihydrokaempferol (DHK) metabolism as well as the metabolism of other substrates such as dihydroquercetin (DHQ), naringenin and eriodictyol. Even more particularly, the present invention provides a genetic sequence encoding a polypeptide having F3'5'H activity when expressed in rose or gerbera or botanically related plants. The  
15 instant invention further relates to antisense and sense molecules or RNAi-inducing molecules corresponding to all or part of the subject genetic sequence or a transcript thereof as well as to genetically modified plants as well as cut flowers, parts and reproductive tissue from such plants. The present invention further relates to promoters which operate efficiently in plants such as rose, gerbera or botanically related plants.

20

#### DESCRIPTION OF PRIOR ART

Reference to any prior art in this specification is not, and should not be taken as, an acknowledgment or any form of suggestion that this prior art forms part of the common  
25 general knowledge in any country.

Bibliographic details of references provided in the subject specification are listed at the end of the specification.

- 2 -

The flower or ornamental plant industry strives to develop new and different varieties of flowers and/or plants. An effective way to create such novel varieties is through the manipulation of flower color. Classical breeding techniques have been used with some success to produce a wide range of colors for almost all of the commercial varieties of flowers and/or plants available today. This approach has been limited, however, by the constraints of a particular species' gene pool and for this reason it is rare for a single species to have the full spectrum of colored varieties. For example, the development of novel colored varieties of plants or plant parts such as flowers, foliage and stems would offer a significant opportunity in both the cut flower and ornamental markets. In the flower or ornamental plant industry, the development of novel colored varieties of major flowering species such as rose, chrysanthemum, tulip, lily, carnation, gerbera, orchid, lisianthus, begonia, torenia, geranium, petunia, nierenbergia, pelargonium, iris, impatiens and cyclamen would be of great interest. A more specific example would be the development of a blue rose or gerbera for the cut flower market.

In addition, the development of novel colored varieties of plant parts such as vegetables, fruits and seeds would offer significant opportunities in agriculture. For example, novel colored seeds would be useful as proprietary tags for plants. Furthermore modifications to flavonoids common to berries or fruits including grapes and apples and their juices including wine have the potential to impart altered style characteristics of value to such fruit and byproduct industries.

Flower color is predominantly due to three types of pigment: flavonoids, carotenoids and betalains. Of the three, the flavonoids are the most common and contribute a range of colors from yellow to red to blue. The flavonoid molecules that make the major contribution to flower color are the anthocyanins, which are glycosylated derivatives of cyanidin and its methylated derivative peonidin, *delphinidin* or *delphinidin-based molecules* and its methylated derivatives petunidin and malvidin and pelargonidin. Anthocyanins are localised in the vacuole of the epidermal cells of petals or the vacuole of the sub epidermal cells of leaves.

- 3 -

The flavonoid pigments are secondary metabolites of the phenylpropanoid pathway. The biosynthetic pathway for the flavonoid pigments (flavonoid pathway) is well established, (Holton and Cornish, *Plant Cell* 7: 1071-1083, 1995; Mol *et al.*, *Trends Plant Sci.* 3: 212-217, 1998; Winkel-Shirley, *Plant Physiol.* 126: 485-493, 2001a; and Winkel-Shirley, *Plant Physiol.* 127: 1399-1404, 2001b) and is shown in Figures 1A and B. Three reactions and enzymes are involved in the conversion of phenylalanine to *p*-coumaroyl-CoA, one of the first key substrates in the flavonoid pathway. The enzymes are phenylalanine ammonia-lyase (PAL), cinnamate 4-hydroxylase (C4H) and 4-coumarate: CoA ligase (4CL). The first committed step in the pathway involves the condensation of three molecules of malonyl-CoA (provided by the action of acetyl CoA carboxylase (ACC) on acetyl CoA and CO<sub>2</sub>) with one molecule of *p*-coumaroyl-CoA. This reaction is catalysed by the enzyme chalcone synthase (CHS). The product of this reaction, 2',4',6', tetrahydroxy-chalcone, is normally rapidly isomerized by the enzyme chalcone flavanone isomerase (CHI) to produce naringenin. Naringenin is subsequently hydroxylated at the 3 position of the central ring by flavanone 3-hydroxylase (F3H) to produce dihydrokaempferol (DHK).

The pattern of hydroxylation of the B-ring of dihydrokaempferol (DHK) plays a key role in determining petal color. The B-ring can be hydroxylated at either the 3', or both the 3' and 5' positions, to produce dihydroquercetin (DHQ) or dihydromyricetin (DHM), respectively. Two key enzymes involved in this part of the pathway are flavonoid 3'-hydroxylase and flavonoid 3', 5'-hydroxylase, both of the cytochrome P450 class of enzymes. Cytochrome P450 enzymes are widespread in nature and genes have been isolated and sequenced from vertebrates, insects, yeasts, fungi, bacteria and plants.

Flavonoid 3'-hydroxylase (F3'H) is a key enzyme in the flavonoid pathway leading to the cyanidin- based pigments which, in many plant species (for example *Rosa spp.*, *Dianthus spp.*, *Petunia spp.*, begonia, cyclamen, impatiens, morning glory and chrysanthemum), contribute to red and pink flower color.

Flavonoid 3', 5'-hydroxylase (F3'5'H) is a key enzyme in the flavonoid pathway leading to the delphinidin- based pigments which, in many plant species (for example, *Petunia spp.*,

- 4 -

*Viola spp.*, *Lisianthus spp.*, *Gentiana spp.*, *Solhya spp.*, *Salvia spp.*, *Clitoria spp.*, *Kennedia spp.*, *Campanula spp.*, *Lavandula spp.*, *Verbena spp.*, *Torenia spp.*, *Delphinium spp.*, *Solanum spp.*, *Cineraria spp.*, *Vitis spp.*, *Bablana stricla*, *Pinus spp.*, *Plcea spp.*, *Larix spp.*, *Phaseolus spp.*, *Vaccinium spp.*, *Cyclamen spp.*, *Iris spp.*, *Pelargonium sp.*,  
 5 *Liparieae*, *Geranium spp.*, *Pisum spp.*, *Lathyrus spp.*, *Catharanthus spp.*, *Malvia spp.*, *Mucuna spp.*, *Vicia spp.*, *Saintpaulia spp.*, *Lagerstroemia spp.*, *bouchina spp.*, *Plumbago spp.*, *Hypocalyptus spp.*, *Rhododendron spp.*, *Linum spp.*, *Macropitilium spp.*, *Hibiscus spp.*, *Hydrangea spp.*, *Cymbidium spp.*, *Millettia spp.*, *Hedysarum spp.*, *Lespedeza spp.*,  
 10 *Asparagus spp.* *Antigonon spp.*, *Pisum spp.*, *Freesia spp.*, *Brunella spp.*, *Clarkia spp.*, etc.), contribute to purple and blue flower color. Many plant species such as roses, gerberas, chrysanthemums and carnations, do not produce delphinidin-based pigments because they lack a F3'5'H activity.

The next step in the pathway, leading to the production of the colored anthocyanins from  
 15 the dihydroflavonols (DHK, DHQ, DHM), involves dihydroflavonol-4-reductase (DFR) leading to the production of the leucoanthocyanidins. The leucoanthocyanidins are subsequently converted to the anthocyanidins, pelargonidin, cyanidin and *delphinidin or delphinidin-based molecules*. These flavonoid molecules are unstable under normal physiological conditions and glycosylation at the 3-position, through the action of  
 20 glycosyltransferases, stabilizes the anthocyanidin molecule thus allowing accumulation of the anthocyanins. In general, the glycosyltransferases transfer the sugar moieties from UDP sugars to the flavonoid molecules and show high specificities for the position of glycosylation and relatively low specificities for the acceptor substrates (Seitz and Hinderer, Anthocyanins. In: *Cell Culture and Somatic Cell Genetics of Plants*. Constabel,  
 25 F. and Vasil, L.K. (eds.), Academic Press, New York, USA, 5: 49-76, 1988). Anthocyanins can occur as 3-monosides, 3-biosides and 3-triosides as well as 3, 5-diglycosides and 3, 7-diglycosides associated with the sugars glucose, galactose, rhamnose, arabinose and xylose (Strack and Wray, In: *The Flavonoids - Advances in Research since 1986*. Harborne, J.B. (ed), Chapman and Hall, London, UK, 1-22, 1993).

- 5 -

Glycosyltransferases involved in the stabilisation of the anthocyanidin molecule include UDP glucose: flavonoid 3-glucosyltransferase (3GT), which transfers a glucose moiety from UDP glucose to the 3-*O*-position of the anthocyanidin molecule to produce anthocyanidin 3-*O*-glucoside.

5

In petunia and pansy (amongst others), anthocyanidin 3-*O*-glucoside are generally glycosylated by another glycosyltransferase, UDP rhamnose: anthocyanidin 3-glucoside rhamnosyltransferase (3RT), which adds a rhamnose group to the 3-*O*-bound glucose of the anthocyanin molecule to produce the anthocyanidin 3-rutinosides, and once acylated, can be further modified by UDP: glucose anthocyanin 5 glucosyltransferase (5GT). However, in roses (amongst others), the anthocyanidin 3-*O*-glucosides are generally glycosylated by another glycosyltransferase, UDP: glucose anthocyanin 5 glucosyltransferase (5GT) to produce anthocyanidin 3, 5 diglucosides.

10

15 Many anthocyanidin glycosides exist in the form of acylated derivatives. The acyl groups that modify the anthocyanidin glycosides can be divided into two major classes based upon their structure. The aliphatic acyl groups include malonic acid or succinic acid and the aromatic class include the hydroxy cinnamic acids such as *p*-coumaric acid, caffeic acid and ferulic acid and the benzoic acids such as *p*-hydroxybenzoic acid.

20

Methylation at the 3' and 5' positions of the B-ring of anthocyanidin glycosides can also occur. Methylation of cyanidin-based pigments leads to the production of peonidin. Methylation of the 3' position of delphinidin-based pigments results in the production of petunidin, whilst methylation of the 3' and 5' positions results in malvidin production.

25

Methylation of malvidin can also occur at the 5-*O* and 7-*O* positions to produce capensinin (5-*O*-methyl malvidin) and 5, 7-di-*O*-methyl malvidin.

30

In addition to the above modifications, pH of the vacuole or compartment where pigments are localised and copigmentation with other flavonoids such as flavonols and flavones can affect petal color. Flavonols and flavones can also be aromatically acylated (Brouillard and

- 6 -

Dangles, In: *The Flavonoids -Advances in Research since 1986*. Harborne, J.B. (ed), Chapman and Hall, London, UK, 1-22, 1993).

- 5 The ability to control F3'5'H activity, or other enzymes involved in the flavonoid pathway, in flowering plants would provide a means of manipulating the color of plant parts such as petals, fruit, leaves, sepals, seeds etc. Different colored versions of a single cultivar could thereby be generated and in some instances a single species would be able to produce a broader spectrum of colors.
- 10 Two nucleotide sequences (referred to herein as SEQ ID NO:1 and SEQ ID NO:3) encoding petunia F3'5'Hs have been cloned (see International Patent Application No. PCT/AU92/00334 and Holton *et al.*, *Nature*, 366: 276-279, 1993a). These sequences were efficient in modulating 3', 5' hydroxylation of flavonoids in petunia (see International Patent Application No. PCT/AU92/00334 incorporated herein by reference and Holton *et*
- 15 *al.*, 1993a, *supra*), tobacco (see International Patent Application No. PCT/AU92/00334 incorporated herein by reference) and carnations (see International Patent Application No. PCT/AU96/00296 incorporated herein by reference). Surprisingly, however, inclusion of these sequences in standard expression cassettes, did not lead to the production of intact or full-length transcripts as detectable by RNA or Northern blot analysis and consequently 3',
- 20 5'-hydroxylated flavonoids were not produced in roses. There is a need, therefore, to identify further genetic sequences encoding F3'5'Hs which efficiently accumulate and are then able to modulate 3', 5' hydroxylation of flavonoids such as anthocyanins in roses and other key commercial plant species.

- 7 -

## SUMMARY OF THE INVENTION

Throughout this specification, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the  
5 inclusion of a stated element or integer or group of elements or integers but not the exclusion of any other element or integer or group of elements or integers.

Nucleotide and amino acid sequences are referred to by a sequence identifier number (SEQ ID NO:). The SEQ ID NOs: correspond numerically to the sequence identifiers <400>1  
10 (SEQ ID NO:1), <400>2 (SEQ ID NO:2), etc.

Genetic sequences encoding a F3'5'H have been identified and cloned from a number of plant species. The F3'5'H genetic sequences when expressed in rose petal tissue results in detectable level of delphinidin or delphinidin-based molecules as determined by a  
15 chromatographic technique such as thin layer chromatography (TLC) or high performance liquid chromatography (HPLC). Alternatively, or in addition, expression of the genetic sequences in rose petal tissue results in a sufficient level and length of transcript which is capable of being translated to F3'5'H. This is conveniently measured as delphinidin or delphinidin-based molecules, detectable using a chromatographic technique such as TLC  
20 or HPLC. The genetic sequences of the present invention permit the modulation of expression of genes encoding this enzyme by, for example, *de novo* expression, over-expression, suppression, antisense inhibition, ribozyme activity, RNAi-induction or methylation-induction. The ability to control F3'5'H synthesis in plants and more specifically in roses or gerberas permits modulation of the composition of individual  
25 anthocyanins as well as alteration of relative levels of flavonols and anthocyanins, thereby enabling the manipulation of color of tissues and/or organs of plants such as petals, leaves, seeds, sepals, fruits etc.

Accordingly, one aspect of the present invention provides an isolated nucleic acid  
30 molecule comprising a sequence of nucleotides encoding or complementary to a sequence encoding a flavonoid 3', 5' hydroxylase (F3'5'H) or a polypeptide having F3'5'H activity

- 8 -

wherein expression of said nucleic acid molecule in a rose petal tissue results in detectable levels of delphinidin or delphinidin-based molecules as measured by a chromatographic technique.

- 5 Another aspect of the present invention is directed to an isolated nucleic acid molecule comprising a sequence of nucleotides encoding or complementary to a sequence encoding a F3'5'H or a polypeptide having F3'5'H activity wherein expression of said nucleic acid molecule in a rose petal tissue results in a sufficient level and length of transcript which is translated to said F3'5'H as determined by detectable levels of delphinidin or delphinidin-  
10 based molecules as measured by a chromatographic technique.

The isolated nucleic acid molecule of the present invention, therefore, encodes a F3'5'H which is capable of more efficient conversion of DHK to DHM in roses than is the F3'5'H encoded by the nucleotide sequence set forth in SEQ ID NO:1 and SEQ ID NO:3 as  
15 measured by delphinidin production in rose petals.

Efficiency as used herein relates to the capability of the F3'5'H enzyme to convert its substrate DHK or DHQ into DHM in a rose cell (or any cell of a commercially important plant such as gerbera). This conversion provides the plant with a substrate (DHM) for other  
20 enzymes of the flavonoid pathway which are present in the plant to further modify the substrate. This modification may include for example, glycosylation, acylation, rhamnosylation and/or methylation, to produce various anthocyanins which contribute to the production of a range of colors. The modulation of 3',5'-hydroxylated anthocyanins in rose is thereby enabled. Efficiency is conveniently assessed by one or more parameters  
25 selected from: extent of F3'5'H transcription, as determined by the amount of intact F3'5'H mRNA produced (as detected by Northern blot analysis); extent of translation of the F3'5'H mRNA, as determined by the amount of translation product produced; extent of F3'5'H enzyme activity as determined by the production of anthocyanin derivatives of DHQ or DHM including delphinidin or delphinidin-based pigments (as detected by TLC or HPLC);  
30 the extent of effect on flower color.



- 9 -

It has also been surprisingly determined that certain combinations of promoter and *F3'5'H* gene sequences that were functional in carnation and petunia were not functional in rose. Surprisingly, only a particular subset of promoter and *F3'5'H* gene sequence combinations resulted in 3', 5'-hydroxylated flavonoids in rose flowers. These included *F3'5'H* sequences isolated from *Viola spp.*, *Salvia spp.*, *Lavandula spp.* and *Sollya spp.* Furthermore, the *Viola F3'5'H* (or pansy *F3'5'H*) sequences were found to result in the highest accumulation of 3', 5'-hydroxylated flavonoids in rose. The novel promoter and *F3'5'H* gene sequence combinations can be employed *inter alia* to modulate the color or flavour or other characteristics of plants or plant parts such as but not limited to flowers, fruits, nuts, roots, stems, leaves or seeds. Thus, the present invention represents a new approach to developing plant varieties having altered color characteristics. Other uses include, for example, the production of novel extracts of *F3'5'H* transformed plants wherein the extract has use, for example, as a flavouring or food additive or health product or beverage or juice or coloring. Beverages may include but are not limited to wines, spirits, teas, coffee, milk and dairy products.

In a preferred embodiment, therefore, the present invention provides an isolated nucleic acid molecule comprising a sequence of nucleotides encoding, or complementary to a sequence encoding pansy *F3'5'H*, salvia *F3'5'H*, lavender *F3'5'H*, kennedia *F3'5'H* or sollya *F3'5'H* or a functional derivative of the enzyme.

The nucleotide sequences encoding the pansy *F3'5'H* (SEQ ID NOs:9 and 11), salvia *F3'5'H* (SEQ ID NOs:13 and 15), sollya *F3'5'H* (SEQ ID NO:17), lavender *F3'5'H* (SEQ ID NO:31) and kennedia *F3'5'H* (SEQ ID NO:26) are defined by sequence identifiers indicated in parentheses. A summary of the sequence identifiers is shown in Table 1.

Accordingly, another aspect of the present invention provides a nucleic acid molecule comprising a nucleotide sequence or complementary nucleotide sequence substantially as set forth in SEQ ID NO:9 (pansy) or SEQ ID NO:11 (pansy) or SEQ ID NO:13 (salvia) or SEQ ID NO:15 (salvia) or SEQ ID NO:17 (sollya) or SEQ ID NO:31 (lavender) or SEQ ID NO:26 (kennedia) or having at least about 50% similarity thereto or capable of

- 10 -

hybridizing to the sequence set forth in SEQ ID NO:9 or SEQ ID NO:11 or SEQ ID NO:13 or SEQ ID NO:15 or SEQ ID NO:17 or SEQ ID NO:31 or SEQ ID NO:26 under low stringency conditions.

- 5 The amino acid sequences of the preferred F3'5'H enzymes are set forth in SEQ ID NO:10 (pansy) or SEQ ID NO:12 (pansy) or SEQ ID NO:14 (salvia) or SEQ ID NO:16 (salvia) or SEQ ID NO:18 (sollya) or SEQ ID NO:32 (lavender) or SEQ ID NO:27 (kennedia).

- 10 A further aspect of the present invention provides a method for producing a transgenic flowering plant capable of synthesizing a F3'5'H said method comprising stably transforming a cell of a suitable plant with a nucleic acid sequence which comprises a sequence of nucleotides encoding said F3'5'H under conditions permitting the eventual expression of said nucleic acid sequence, regenerating a transgenic plant from the cell and growing said transgenic plant for a time and under conditions sufficient to permit the
- 15 expression of the nucleic acid sequence. The expression of the nucleic acid sequence generally results in a transcription of sufficient level and length to encode a F3'5'H. This is conveniently determined by detectable levels of delphinidin or delphinidin-based molecules as measured by chromatographic techniques such as TLC or HPLC. The transgenic plant may thereby produce a non-indigenous F3'5'H at elevated levels relative to
- 20 the amount expressed in a comparable non-transgenic plant. This generally results in a visually detectable color change in the plant or plant part or preferably in the inflorescence or flowers of said plant.

- 25 Another aspect of the present invention contemplates a method for producing a transgenic plant with reduced F3'5'H activity, said method comprising stably transforming a cell of a suitable plant with a nucleic acid molecule which comprises a sequence of nucleotides encoding or complementary to a sequence encoding a F3'5'H activity, regenerating a transgenic plant from the cell and where necessary growing said transgenic plant under conditions sufficient to permit the expression of the nucleic acid.

30

- 11 -

Yet another aspect of the present invention contemplates a method for producing a genetically modified plant with reduced F3'5'H activity, said method comprising altering the F3'5'H gene through modification of the indigenous sequences *via* homologous recombination from an appropriately altered *F3'5'H* gene or derivative or part thereof introduced into the plant cell, and regenerating the genetically modified plant from the cell.

Still another aspect of the present invention contemplates a method for producing a transgenic flowering plant exhibiting altered floral or inflorescence properties, said method comprising stably transforming a cell of a suitable plant with a nucleic acid sequence of the present invention, regenerating a transgenic plant from the cell and growing said transgenic plant for a time and under conditions sufficient to permit the expression of the nucleic acid sequence.

Still a further aspect of the present invention contemplates a method for producing a transgenic flowering plant exhibiting altered floral or inflorescence properties, said method comprising alteration of the *F3'5'H* gene through modification of nucleotide sequences *via* homologous recombination from an appropriately altered *F3'5'H* gene or derivative or part thereof introduced into the plant cell, and regenerating the genetically modified plant from the cell.

Even yet another aspect of the present invention extends to a method for producing a transgenic plant capable of expressing a recombinant gene encoding a F3'5'H or part thereof or which carries a nucleic acid sequence which is substantially complementary to all or a part of a mRNA molecule encoding said F3'5'H, said method comprising stably transforming a cell of a suitable plant with the isolated nucleic acid molecule comprising a sequence of nucleotides encoding, or complementary to a sequence encoding, a F3'5'H, where necessary under conditions permitting the eventual expression of said isolated nucleic acid molecule, and regenerating a transgenic plant from the cell.

- 12 -

Even still another aspect of the present invention extends to all transgenic plants or parts of transgenic plants or progeny of the transgenic plants containing all or part of the nucleic acid sequences of the present invention, or antisense forms thereof and/or any homologs or related forms thereof and, in particular, those transgenic plants which exhibit altered floral or inflorescence properties.

Even still another aspect of the present invention extends to all transgenic plants or parts of transgenic plants or progeny of the transgenic plants containing all or part of the nucleic acid sequences of the present invention, or antisense forms thereof and/or any homologs or related forms thereof and, in particular, those transgenic plants which exhibit altered aerial parts of the plant such as fruit, berries, sepal, bract, petiole, peduncle, ovaries, anthers or stem properties.

Another aspect of the present invention contemplates the use of the extracts from transgenic plants or plant parts transgenic plants or progeny of the transgenic plants containing all or part of the nucleic acid sequences of the present invention and, in particular, the extracts from those transgenic plants when used as a flavouring or food additive or health product or beverage or juice or coloring.

A further aspect of the present invention is directed to recombinant forms of F3'5'H.

Another aspect of the present invention contemplates the use of the genetic sequences described herein in the manufacture of a genetic construct capable of expressing a F3'5'H or down-regulating an indigenous F3'5'H enzyme in a plant.

Yet another aspect of the present invention is directed to a prokaryotic or eukaryotic organism carrying a genetic sequence encoding a F3'5'H extrachromasomally in plasmid form.

- 13 -

Still another aspect of the present invention extends to a recombinant polypeptide comprising a sequence of amino acids substantially as set forth in SEQ ID NO:10 or SEQ ID NO:12 or SEQ ID NO:14 or SEQ ID NO:16 or SEQ ID NO:18 or SEQ ID NO:32 or SEQ ID NO:27 or an amino acid sequence having at least about 50% similarity to SEQ ID NO:10 or SEQ ID NO:12 or SEQ ID NO:14 or SEQ ID NO:16 or SEQ ID NO:18 or SEQ ID NO:32 or SEQ ID NO:27 or a derivative of said polypeptide.

The present invention further provides promoters which operate efficiently in plants such as rose and gerbera or botanically related plants. Such promoters include a rose CHS promoter, chrysanthemum CHS promoter and a CaMV 35S promoter.

A summary of sequence identifiers used throughout the subject specification is provided in Table 1:

# TABLE 1

## Summary of sequence identifiers

SEQ ID NO:	NAME	SPECIES	TYPE OF SEQ	DESCRIPTION
1	<i>petHf1.nt</i>	<i>Petunia hybrida</i>	nucleotide	<i>F3'5'H cDNA</i>
2	<i>petHf1.aa</i>	<i>Petunia hybrida</i>	amino acid	<i>translation of F3'5'H cDNA</i>
3	<i>petHf2.nt</i>	<i>Petunia hybrida</i>	nucleotide	<i>F3'5'H cDNA</i>
4	<i>petHf2.aa</i>	<i>Petunia hybrida</i>	amino acid	<i>translation of F3'5'H cDNA</i>
5	<i>RoseCHS promoter</i>	<i>Rosa hybrida</i>	nucleotide	<i>promoter fragment</i>
6	D8 oligo#2	<i>Petunia hybrida</i>	nucleotide	<i>oligonucleotide</i>
7	D8 oligo #4	<i>Petunia hybrida</i>	nucleotide	<i>oligonucleotide</i>
8	chrysanCHSATG	chrysanthemum	nucleotide	<i>oligonucleotide</i>
9	<i>BP#18.nt</i>	<i>Viola spp.</i>	nucleotide	<i>F3'5'H cDNA</i>
10	<i>BP#18.aa</i>	<i>Viola spp.</i>	amino acid	<i>translation of F3'5'H cDNA</i>
11	<i>BP#40.nt</i>	<i>Viola spp.</i>	nucleotide	<i>F3'5'H cDNA</i>
12	<i>BP#40.aa</i>	<i>Viola spp.</i>	amino acid	<i>translation of F3'5'H cDNA</i>
13	<i>Sal#2.nt</i>	<i>Salvia spp.</i>	nucleotide	<i>F3'5'H cDNA</i>

- 14 -

SEQ ID NO:	NAME	SPECIES	TYPE OF SEQ	DESCRIPTION
14	Sal#2.aa	Salvia spp.	amino acid	translation of F3'5'H cDNA
15	Sal#47.nt	Salvia spp.	nucleotide	F3'5'H cDNA
16	Sal#47.aa	Salvia spp.	amino acid	translation of F3'5'H cDNA
17	Soll#5.nt	Sollya spp.	nucleotide	F3'5'H cDNA
18	Soll#5.aa	Sollya spp.	amino acid	translation of F3'5'H cDNA
19	FLS-Nco	Petunia hybrida	nucleotide	oligonucleotide
20	BpeaHF2.nt	Clitoria ternatea	nucleotide	F3'5'H cDNA
21	BpeaHF2.aa	Clitoria ternatea	amino acid	translation of F3'5'H cDNA
22	Gen#48.nt	Gentiana triflora	nucleotide	F3'5'H cDNA
23	Gen#48.aa	Gentiana triflora	amino acid	translation of F3'5'H cDNA
24	PetD8 5'	Petunia hybrida	nucleotide	oligonucleotide
25	Bpea primer	Clitoria ternatea	nucleotide	oligonucleotide
26	Kenn#31.nt	Kennedia spp.	nucleotide	F3'5'H cDNA
27	Kenn#31.aa	Kennedia spp.	amino acid	translation of F3'5'H cDNA
28	chrysCHS.nt	chrysanthemum	nucleotide	CHS cDNA
29	chrysCHS.aa	chrysanthemum	amino acid	translation of CHS cDNA
30	chrysCHS promoter	chrysanthemum	nucleotide	promoter fragment
31	LBG.nt	Lavandula nil	nucleotide	F3'5'H cDNA
32	LBG.aa	Lavandula nil	amino acid	translation of F3'5'H cDNA

## BRIEF DESCRIPTION OF THE FIGURES

Figures 1A and 1B are schematic representations of the biosynthesis pathway for the flavonoid pigments. Figure 1A illustrates the general production of the anthocyanidin 3-glucosides that occur in most plants that produce anthocyanins. Figure 1B represents further modifications of anthocyanins that occur in petunia. Enzymes involved in the pathway have been indicated as follows: PAL = Phenylalanine ammonia-lyase; C4H = Cinnamate 4-hydroxylase; 4CL = 4-coumarate: CoA ligase; CHS = Chalcone synthase; CHI = Chalcone flavanone isomerase; F3H = Flavanone 3-hydroxylase; DFR = Dihydroflavonol-4-reductase; ANS = Anthocyanidin synthase, 3GT = UDP-glucose: flavonoid 3-O-glucosyltransferase; 3RT = UDP rhamnose: anthocyanidin 3-glucoside rhamnosyltransferase, AR-AT = Anthocyanidin-rutinoside acyltransferase, 5GT = Anthocyanin 5-glucosyltransferase; 3' OMT = Anthocyanin 3' O-methyltransferase, 3'5' OMT = Anthocyanin 3', 5' O-methyltransferase. Other abbreviations include: DHK = dihydrokaempferol, DHQ = dihydroquercetin, DHM = dihydromyricetin,

TABLE 2: Descriptions of the abbreviations used in Figures 2 to 52

ABBREVIATION	DESCRIPTION
Amp	ampicillin resistance gene which confers resistance to the antibiotic ampicillin
ColE1ori	plasmid origin of replication
fl ori (+)	fl filamentous phage origin of replication
GentR	gentamycin resistance gene which confers resistance to the antibiotic gentamycin
LB	left border of the T-DNA
<i>npIII</i>	the neomycin phosphotransferase III gene which confers resistance to the antibiotic kanamycin
ori pRi	plasmid origin of replication
ori 322	plasmid origin of replication
pACYC ori	modified replicon from pACYC184 from <i>E. coli</i>

- 16 -

pVS1	a broad host range origin of replication from a plasmid from <i>Pseudomonas aeruginosa</i>
rev	approximate location of the M13 reverse primer site used in sequence analysis
RB	right border of the T-DNA
TetR	tetracycline resistance gene which confers resistance to the antibiotic tetracycline
-20	approximate location of the M13 -20 primer site used in sequence analysis
RK2	broad host range Gram-negative plasmid RK2 origin

Figure 2 is a diagrammatic representation of the plasmid pCGP602, pCGP601 and pCGP176 containing petunia *F3'5'H petHf1* cDNA clones from *P. hybrida* cv. OGB. The petunia *F3'5'H petHf1* fragment was used in the preparation of constructs containing the petunia *F3'5'H* cDNA clone. <sup>32</sup>P-labelled fragments of the 1.6 kb *BspHI/FspI* fragment were used to probe petal cDNA libraries. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 3 is a diagrammatic representation of the plasmid pCGP175 containing the petunia *F3'5'H petHf2* cDNA clone from *P. hybrida* cv. OGB. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 4 is a diagrammatic representation of the plasmid pCGP1303 containing a subclone of the petunia *F3'5'H petHf1* cDNA clone from pCGP601. The construction of pCGP1303 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 5 is a diagrammatic representation of the binary plasmid pCGP1452. The *AmCHS* 5': *petHf1*: *petD8* 3' gene from pCGP485 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1452 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.



Figure 6 is a diagrammatic representation of the binary plasmid pWTT2132 (DNAP) containing the 35S 5': SuRB selectable marker gene and a multi-cloning site. A description of pWTT2132 is given in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

5

Figure 7 is a diagrammatic representation of the plasmid pCGP725. The *AmCHS* 5': *petHf1: petD8* 3' gene from pCGP485 was cloned into pBluescript II (KS (+) vector. The construction of pCGP725 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

10

Figure 8 is a diagrammatic representation of the binary plasmid pCGP1453. The *Mac: petHf1: mas* 3' gene from pCGP628 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1453 is described in Example 4. Selected restriction endonuclease sites are marked.

15

Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 9 is a diagrammatic representation of the binary plasmid pCGP1457. The *petD8* 5': *petHf1: petD8* 3' gene from pCGP1107 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1457 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

20

Figure 10 is a diagrammatic representation of the binary plasmid pCGP1461. The *shortpetFLS* 5': *petHf1: petFLS* 3' gene from pCGP497 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1461 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

25

Figure 11 is a diagrammatic representation of the binary plasmid pCGP1616. The *petRT* 5': *petHf1: nos* 3' gene from pCGP846 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimaeric *SuRB* gene. The construction of

30

pCGP1616 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 12 is a diagrammatic representation of the binary plasmid pCGP1623. The *mas/35S: petHf1: ocs 3'* gene from pCGP1619 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1623 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 13 is a diagrammatic representation of the binary plasmid pCGP1638. The *CaMV 35S: petHf1: nos 3'* gene from pCGP1636 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1636 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 14 is a diagrammatic representation of the binary plasmid pCGP1860. The *RoseCHS 5': petHf1: nos 3'* gene from pCGP200 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1860 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 15 is a diagrammatic representation of the binary plasmid pCGP2123. The *CaMV35S: petHf2: ocs 3'* gene from pCGP2109 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2123 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 16 is a diagrammatic representation of the binary plasmid pCGP1988. The multi-cloning site of the binary vector pWTT2132 (DNAP) was replaced with the multi-cloning site from pNEB193 (New England Biolabs). The construction of pCGP1988 is described in

- 19 -

Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 17 is a diagrammatic representation of the plasmid pCGP2105. The 35S 5': *ocs* 3' expression cassette with multiple restriction endonuclease sites between the promoter and terminator fragments is in a pBluescript SK (+) vector backbone. The construction of pCGP2105 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 18 is a diagrammatic representation of the binary plasmid pCGP1307. The *petD8* 5': *GUS*: *petD8* 3' gene from pCGP1106 was cloned into the binary vector pCGN1548 in a tandem orientation to the chimaeric *nptII* selectable marker gene. The construction of pCGP1307 is described in Example 6. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 19 is a diagrammatic representation of the binary plasmid pCGP1506. The *longpetFLS* 5': *GUS*: *petFLS* 3' gene from pCGP496 was cloned into the binary vector pBIN19 in a tandem orientation to the chimaeric *nptII* selectable marker gene. The construction of pCGP1506 is described in Example 6. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 20 is a diagrammatic representation of the binary plasmid pCGP1626. The *ChrysCHS* 5': *GUS*: *nos* 3' gene from pCGP1622 was cloned into the binary vector pWTT2132 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1626 is described in Example 6. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 21 is a diagrammatic representation of the binary plasmid pCGP1641. The *petRT* 5': *GUS*: *petRT* 3' gene from pCGP1628 was cloned into the binary vector pWTT2132 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1641 is

- 20 -

described in Example 6. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

5 **Figure 22** is a diagrammatic representation of the binary plasmid pCGP1861. The *RoseCHS* 5': *GUS: nos* 3' gene from pCGP197 was cloned into the binary vector pWTT2132 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1861 is described in Example 6. Refer to Table 2 and Table 4 for a description of the abbreviations.

10 **Figure 23** is a diagrammatic representation of the binary plasmid pCGP1953. The *AmCHS* 5': *GUS: petD8* 3' gene from pCGP1952 was cloned into the binary vector pWTT2132 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1953 is described in Example 6. Refer to Table 2 and Table 4 for a description of the abbreviations.

15 **Figure 24** is a diagrammatic representation of the binary plasmid pWTT2084 (DNAP) containing a 35S 5': *GUS: ocs* 3' gene in a convergent orientation to the chimaeric *SuRB* selectable marker gene. A description of pWTT2084 is given in Example 6. Refer to Table 2 and Table 4 for a description of the abbreviations.

20 **Figure 25** is a diagrammatic representation of the plasmid pCGP1959 containing the *F3'5'H BP#18* cDNA clone from *Viola spp.* cv Black Pansy in a pBluescript SK II (+) backbone. A description of pCGP1959 is given in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

25 **Figure 26** is a diagrammatic representation of the plasmid pCGP1961 containing the *F3'5'H BP#40* cDNA clone from *Viola spp.* cv Black Pansy in a pBluescript SK II (+) backbone. A description of pCGP1961 is given in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

30

- 21 -

Figure 27 is a diagrammatic representation of the binary plasmid pCGP1972. The *AmCHS* 5': BP#18: *petD8* 3' gene from pCGP1970 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1972 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 28 is a diagrammatic representation of the binary plasmid pCGP1973. The *AmCHS* 5': BP#40: *petD8* 3' gene from pCGP1971 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1973 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 29 is a diagrammatic representation of the binary plasmid pCGP1967. The *CaMV* 35S: BP#18: *ocs* 3' gene from pCGP1965 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1967 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 30 is a diagrammatic representation of the binary plasmid pCGP1969. The *CaMV* 35S: BP#40: *ocs* 3' gene from pCGP1966 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1969 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 31 is a diagrammatic representation of the plasmid pCGP1995 containing the *F3'5'H Sal#2* cDNA clone from *Salvia spp.* in a pBluescript SK II (+) backbone. A description of pCGP1995 is given in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

- 22 -

Figure 32 is a diagrammatic representation of the plasmid pCGP1999 containing the *F3'5'H Sal#47* cDNA clone from *Salvia spp* in a pBluescript SK II (+) backbone. A description of pCGP1999 is given in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

5

Figure 33 is a diagrammatic representation of the binary plasmid pCGP2121. The *AmCHS* 5': *Sal#2: petD8* 3' gene from pCGP2116 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2121 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

10

Figure 34 is a diagrammatic representation of the binary plasmid pCGP2122. The *AmCHS* 5': *Sal#47: petD8* 3' gene from pCGP2117 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2122 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

15

Figure 35 is a diagrammatic representation of the binary plasmid pCGP2120. The *CaMV* 35S:*Sal#2:ocs* 3' gene from pCGP2112 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2120 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

20

Figure 36 is a diagrammatic representation of the binary plasmid pCGP2119. The *CaMV* 35S:*Sal#47:ocs* 3' gene from pCGP2111 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2119 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

25

- 23 -

Figure 37 is a diagrammatic representation of the plasmid pCGP2110 containing the *F3'5'H Soll#5* cDNA clone from *Sollya spp.* in a pBluescript SK II (+) backbone. A description of pCGP2110 is given in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

5

Figure 38 is a diagrammatic representation of the binary plasmid pCGP2130. The *AmCHS 5': Soll#5: petD8 3'* gene from pCGP2128 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2130 is described in Example 7. Refer to Table 2 and Table 4 for a description of the abbreviations.

10

Figure 39 is a diagrammatic representation of the binary plasmid pCGP2131. The *CaMV 35S: Soll#5:ocs 3'* gene from pCGP2129 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2131 is described in Example 7. Refer to Table 2 and Table 4 for a description of the abbreviations.

15

Figure 40 is a diagrammatic representation of the plasmid pCGP2231 containing the *F3'5'H Kenn#31* cDNA clone from *Kennedia spp.* in a pBluescript SK II (+) backbone. A description of pCGP2231 is given in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

20

Figure 41 is a diagrammatic representation of the binary plasmid pCGP2256. The *AmCHS 5': Kenn#31: petD8 3'* gene from pCGP2242 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2256 is described in Example 7. Refer to Table 2 and Table 4 for a description of the abbreviations.

25

- 24 -

Figure 42 is a diagrammatic representation of the binary plasmid pCGP2252. The *CaMV* 35S: *Kenn#31:ocs* 3' gene from pCGP2236 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2252 is described in Example 7. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 43 is a diagrammatic representation of the plasmid pBHF2F containing the full-length *F3'5'H BpeaHF2* cDNA clone from *Clitoria ternatea* in a pBluescript SK II (+) backbone. A description of pBHF2F is given in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 44 is a diagrammatic representation of the binary plasmid pCGP2135. The *AmCHS* 5': *BpeaHF2: petD8* 3' gene from pCGP2133 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2135 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 45 is a diagrammatic representation of the binary plasmid pBEBF5. The *eCaMV* 35S: *BpeaHF2: nos* 3' gene was constructed by replacing the *GUS* fragment from pBE2113-GUSs with the *Clitoria F3'5'H BpeaHF2* cDNA clone from pBHF2F. The construction of pBEBF5 is described in Example 7. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 46 is a diagrammatic representation of the binary plasmid pCGP2134. The *CaMV* 35S: *BpeaHF2: ocs* 3' gene from pCGP2132 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2134 is described in Example 7. Refer to Table 2 and Table 4 for a description of the abbreviations.

30



- 25 -

Figure 47 is a diagrammatic representation of the plasmid pG48 containing the *F3'5'H Gen#48* cDNA clone from *Gentiana triflora* in a pBluescript SK II (+) backbone. A description of pG48 is given in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

5

Figure 48 is a diagrammatic representation of the binary plasmid pCGP1498. The *AmCHS 5': Gen#48: petD8 3'* gene from pCGP1496 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1498 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

10

Figure 49 is a diagrammatic representation of the binary plasmid pBEGHF48. The *eCaMV 35S: Gen#48: nos 3'* gene was constructed by replacing the *GUS* fragment from pBE2113-GUSs with the *Gentiana F3'5'H Gen#48* cDNA clone from pG48. The construction of pBEGHF48 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

15

Figure 50 is a diagrammatic representation of the binary plasmid pCGP1982. The *CaMV 35S: Gen#48: ocs 3'* gene from pCGP1981 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1982 is described in Example 7. Refer to Table 2 and Table 4 for a description of the abbreviations.

20

Figure 51 is a diagrammatic representation of the plasmid pLFH8 containing the *F3'5'H LBG* cDNA clone from *Lavandula nil* in a pBluescript SK II (+) backbone. A description of pLFH8 is given in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

25

- 26 -

Figure 52 is a diagrammatic representation of the binary plasmid pBELF8. The *eCaMV* 35S: *LBG: nos* 3' gene was constructed by replacing the *GUS* fragment from pBE2113-*GUS*s with the *Lavandula F3'5'H LBG* cDNA clone from pLHF8. The construction of pBELF8 is described in Example 7. Refer to Table 2 and Table 4 for a description of the

5 abbreviations.

## DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

In accordance with the present invention, genetic sequences encoding polypeptides having F3'5'H activity have been identified, cloned and assessed. The recombinant genetic sequences of the present invention permit the modulation of expression of genes encoding this enzyme by, for example, *de novo* expression, over-expression, sense suppression, antisense inhibition, ribozyme, minizyme and DNAzyme activity, RNAi-induction or methylation-induction or other transcriptional or post-transcriptional silencing activities. RNAi-induction includes genetic molecules such as hairpin, short double stranded DNA or RNA, and partially double stranded DNAs or RNAs with one or two single stranded nucleotide over hangs. The ability to control F3'5'H synthesis in plants permits modulation of the composition of individual anthocyanins as well as alteration of relative levels of flavonols and anthocyanins, thereby enabling the manipulation of petal color. Moreover, the present invention extends to plants and reproductive or vegetative parts thereof including flowers, fruits, seeds, vegetables, leaves, stems and the like. The present invention further extends to ornamental transgenic or genetically modified plants. The term "transgenic" also includes progeny plants and plants from subsequent genetics and/or crosses thereof from the primary transgenic plants.

Accordingly, one aspect of the present invention provides an isolated nucleic acid molecule comprising a sequence of nucleotides encoding or complementary to a sequence encoding a F3'5'H or a polypeptide having F3'5'H activity wherein expression of said nucleic acid molecule in a rose petal tissue results in detectable levels of delphinidin or delphinidin-based molecules as measured by a chromatographic technique.

Another aspect of the present invention is directed to an isolated nucleic acid molecule comprising a sequence of nucleotides encoding or complementary to a sequence encoding a F3'5'H or a polypeptide having F3'5'H activity wherein expression of said nucleic acid molecule in a rose petal tissue results in a sufficient level and length of transcript which is translated to said F3'5'H as determined by detectable levels of delphinidin or delphinidin-based molecules as measured by a chromatographic technique.

- 28 -

A further aspect of the present invention is directed to an isolated nucleic acid molecule comprising a sequence of nucleotides encoding or complementary to a sequence encoding a F3'5'H or a polypeptide having F3'5'H activity wherein expression of said nucleic acid molecule in a rose petal tissue results a full-length transcript which is detectable by Northern blot analysis of total RNA isolated from rose petals.

The present invention is described and exemplified herein by reference to the identification, cloning and manipulation of genetic sequences encoding a F3'5'H which acts on DHK as well as DHQ. Preferably, the F3'5'H enzyme is a pansy, salvia, sollya lavender or kennedia F3'5'H. The F3'5'H enzyme may also be considered to include a polypeptide or protein having a F3'5'H activity or F3'5'H-like activity. The latter encompasses derivatives having altered F3'5'H activities.

A preferred aspect of the present invention, therefore, is directed to an isolated nucleic acid molecule comprising a sequence of nucleotides encoding, or complementary to a sequence encoding a F3'5'H or a functional mutant, derivative, part, fragment, homolog or analog thereof wherein the nucleic acid molecule is characterized by the following:

(i) the F3'5'H transcript in rose petal tissue is of sufficient level and size to encode a F3'5'H resulting in detectable delphinidin or delphinidin-based molecules in the rose petal tissue as measured by a chromatographic procedure (eg. TLC or HPLC);

(ii) the F3'5'H transcript in rose petal tissue is full-length and detected by Northern blot analysis of total RNA isolated from rose petal tissue

(iii) the F3'5'H in rose petal tissue results in detectable delphinidin or delphinidin-based molecules as measured by a chromatographic procedure (eg. TLC or HPLC); and/or

(iv) the F3'5'H results in a visual color change in rose petal tissue.

- 29 -

The term delphinidin-based pigments includes the anthocyanidin, delphinidin or any derivatives thereof including but not limited to glycosylated, acylated, methylated or other modified forms. Methylated forms of delphinidin include but are not limited to the anthocyanidin petunidin (methylated at the 3'-position), malvidin (methylated at the 3' and 5' position), 5-O methyl malvidin (methylated at the 5, 3' and 5' positions), 5, 7-O dimethyl malvidin (methylated at the 5, 7, 3' and 5' positions). The methylated anthocyanidins can also be modified by glycosylation and acylation. The term anthocyanins defines glycosylated forms of the respective anthocyanidins.

10 By the term "nucleic acid molecule" is meant a genetic sequence in a non-naturally occurring condition. Generally, this means isolated away from its natural state or synthesized or derived in a non-naturally-occurring environment. More specifically, it includes nucleic acid molecules formed or maintained *in vitro*, including genomic DNA fragments recombinant or synthetic molecules and nucleic acids in combination with  
15 heterologous nucleic acids. It also extends to the genomic DNA or cDNA or part thereof encoding F3'5'H or a part thereof in reverse orientation relative to its own or another promoter. It further extends to naturally occurring sequences following at least a partial purification relative to other nucleic acid sequences.

20 The term "genetic sequences" is used herein in its most general sense and encompasses any contiguous series of nucleotide bases specifying directly, or *via* a complementary series of bases, a sequence of amino acids in a F3'5'H enzyme. Such a sequence of amino acids may constitute a full-length F3'5'H such as is set forth in SEQ ID NO: 10 (pansy) or SEQ ID NO:12 (pansy) or SEQ ID NO:14 (salvia) or SEQ ID NO:16 (salvia) or SEQ ID  
25 NO:18 (sollya) or SEQ ID NO:32 (lavender) or SEQ ID NO:27 (kennedia) or an active truncated form thereof or may correspond to a particular region such as an N-terminal, C-terminal or internal portion of the enzyme. A genetic sequence may also be referred to as a sequence of nucleotides or a nucleotide sequence and includes a recombinant fusion of two or more sequences.

30

- 30 -

In accordance with the above aspects of the present invention there is provided a nucleic acid molecule comprising a nucleotide sequence or complementary nucleotide sequence substantially as set forth in SEQ ID NO:9 (pansy) or SEQ ID NO:11 (pansy) or SEQ ID NO:13 (salvia) or SEQ ID NO:15 (salvia) or SEQ ID NO:17 (sollya) or SEQ ID NO:31 (lavender) or SEQ ID NO:26 (kennedia) or having at least about 50% similarity thereto or capable of hybridizing to the sequence set forth in SEQ ID NO:9 or SEQ ID NO:11 or SEQ ID NO:13 or SEQ ID NO:15 or SEQ ID NO:17 or SEQ ID NO:31 or SEQ ID NO:26 under low stringency conditions.

Table 1 provides a summary of the sequence identifiers.

Alternative percentage similarities and identities (at the nucleotide or amino acid level) encompassed by the present invention include at least about 60% or at least about 65% or at least about 70% or at least about 75% or at least about 80% or at least about 85% or at least about 90% or above, such as about 95% or about 96% or about 97% or about 98% or about 99%, such as at least about 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 100%.

In a particularly preferred embodiment, there is provided an isolated nucleic acid molecule comprising a nucleotide sequence or complementary nucleotide sequence substantially as set forth in SEQ ID NO:9 (pansy) or SEQ ID NO:11 (pansy) or SEQ ID NO:13 (salvia) or SEQ ID NO:15 (salvia) or SEQ ID NO:17 (sollya) or SEQ ID NO:31 (lavender) or SEQ ID NO:26 (kennedia) or having at least about 50% similarity thereto or capable of hybridizing to the sequence set forth in SEQ ID NO:1 (petunia) or SEQ ID NO:3 (petunia) or complementary strands of either under low stringency conditions, wherein said nucleotide sequence encodes a polypeptide having a F3'5'H activity.

For the purposes of determining the level of stringency to define nucleic acid molecules capable of hybridizing to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:9 or SEQ ID NO:11

- 31 -

or SEQ ID NO:13 or SEQ ID NO:15 or SEQ ID NO:17 or SEQ ID NO:31 or SEQ ID NO:26 reference herein to a low stringency includes and encompasses from at least about 0% to at least about 15% v/v formamide and from at least about 1M to at least about 2 M salt for hybridization, and at least about 1 M to at least about 2 M salt for washing conditions. Generally, low stringency is from about 25-30°C to about 42°C. The temperature may be altered and higher temperatures used to replace the inclusion of formamide and/or to give alternative stringency conditions. Alternative stringency conditions may be applied where necessary, such as medium stringency, which includes and encompasses from at least about 16% v/v to at least about 30% v/v formamide and from at least about 0.5 M to at least about 0.9 M salt for hybridization, and at least about 0.5 M to at least about 0.9 M salt for washing conditions, or high stringency, which includes and encompasses from at least about 31% v/v to at least about 50% v/v formamide and from at least about 0.01 M to at least about 0.15 M salt for hybridization, and at least about 0.01 M to at least about 0.15 M salt for washing conditions. In general, washing is carried out  $T_m = 69.3 + 0.41 (G+C)\%$  (Marmur and Doty, *J. Mol. Biol.* 5: 109, 1962). However, the  $T_m$  of a duplex DNA decreases by 1°C with every increase of 1% in the number of mismatch base pairs (Bonner and Laskey, *Eur. J. Biochem.* 46: 83, 1974). Formamide is optional in these hybridization conditions. Accordingly, particularly preferred levels of stringency are defined as follows: low stringency is 6 x SSC buffer, 1.0% w/v SDS at 25-42°C; a moderate stringency is 2 x SSC buffer, 1.0% w/v SDS at a temperature in the range 20°C to 65°C; high stringency is 0.1 x SSC buffer, 0.1% w/v SDS at a temperature of at least 65°C.

Another aspect of the present invention provides a nucleic acid molecule comprising a sequence of nucleotides encoding or complementary to a sequence encoding an amino acid sequence substantially as set forth in SEQ ID NO:10 or SEQ ID NO:12 or SEQ ID NO:14 or SEQ ID NO:16 or SEQ ID NO:18 or SEQ ID NO:32 or SEQ ID NO:27 or an amino acid sequence having at least about 50% similarity thereto.

The term similarity as used herein includes exact identity between compared sequences at the nucleotide or amino acid level. Where there is non-identity at the nucleotide level,

- 32 -

similarity includes differences between sequences which result in different amino acids that are nevertheless related to each other at the structural, functional, biochemical and/or conformational levels. Where there is non-identity at the amino acid level, similarity includes amino acids that are nevertheless related to each other at the structural, functional, biochemical and/or conformational levels. In a particularly preferred embodiment, nucleotide and sequence comparisons are made at the level of identity rather than similarity.

Terms used to describe sequence relationships between two or more polynucleotides or polypeptides include "reference sequence", "comparison window", "sequence similarity", "sequence identity", "percentage of sequence similarity", "percentage of sequence identity", "substantially similar" and "substantial identity". A "reference sequence" is at least 12 but frequently 15 to 18 and often at least 25 or above, such as 30 monomer units, inclusive of nucleotides and amino acid residues, in length. Because two polynucleotides may each comprise (1) a sequence (i.e. only a portion of the complete polynucleotide sequence) that is similar between the two polynucleotides, and (2) a sequence that is divergent between the two polynucleotides, sequence comparisons between two (or more) polynucleotides are typically performed by comparing sequences of the two polynucleotides over a "comparison window" to identify and compare local regions of sequence similarity. A "comparison window" refers to a conceptual segment of typically 12 contiguous residues that is compared to a reference sequence. The comparison window may comprise additions or deletions (i.e. gaps) of about 20% or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Optimal alignment of sequences for aligning a comparison window may be conducted by computerized implementations of algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Drive Madison, WI, USA) or by inspection and the best alignment (i.e. resulting in the highest percentage homology over the comparison window) generated by any of the various methods selected. Reference also may be made to the BLAST family of programs as, for example, disclosed by Altschul *et al.* (*Nucl. Acids Res.* 25: 3389-3402, 1997). A detailed discussion of sequence analysis can be found in Unit



- 33 -

19.3 of Ausubel *et al.* ("Current Protocols in Molecular Biology" John Wiley & Sons Inc, 1994-1998, Chapter 15, 1998).

The terms "sequence similarity" and "sequence identity" as used herein refers to the extent that sequences are identical or functionally or structurally similar on a nucleotide-by-nucleotide basis or an amino acid-by-amino acid basis over a window of comparison. Thus, a "percentage of sequence identity", for example, is calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g. A, T, C, G, I) or the identical amino acid residue (e.g. Ala, Pro, Ser, Thr, Gly, Val, Leu, Ile, Phe, Tyr, Trp, Lys, Arg, His, Asp, Glu, Asn, Gln, Cys and Met) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity. For the purposes of the present invention, "sequence identity" will be understood to mean the "match percentage" calculated by the DNASIS computer program (Version 2.5 for windows; available from Hitachi Software engineering Co., Ltd., South San Francisco, California, USA) using standard defaults as used in the reference manual accompanying the software. Similar comments apply in relation to sequence similarity.

The nucleic acid molecules of the present invention may be further characterized by having, or previously having, prior to derivatization on overall lower AT content (or higher GC content) compared to a nucleic acid molecule which encodes a F3'5'H but which does not result in detectable intact transcript in rose petal tissue or, when expressed, does not result in detectable delphinidin or delphinidin-based molecules, as measured by a chromatographic procedure such as TLC or HPLC. Furthermore, the % of A's or T's in the third position of a codon is also lower than other F4'5'H enzymes. Reference herein to a chromatographic procedure includes a related procedure. By "related" means a technically related procedure or a procedure which provides a similar result. Examples of related procedures include other forms of chromatography (eg. gas chromatography).

- 34 -

In addition, nucleotide sequences which do not express well in rose tissue may be modified such as in reducing overall % AT or at least reduce the levels of % AT in the third position of a codon. Such time expression in rose tissue is elevated.

- 5 The nucleic acid sequences contemplated herein also encompass oligonucleotides useful as genetic probes for amplification reactions or as antisense or sense molecules capable of regulating expression of the corresponding gene in a plant. Sense molecules include hairpin constructs, short double stranded DNAs and RNAs and partially double stranded DNAs and RNAs which one or more single stranded nucleotide over hangs. An antisense
- 10 molecule as used herein may also encompass a genetic construct comprising the structural genomic or cDNA gene or part thereof in reverse orientation relative to its or another promoter. It may also encompass a homologous genetic sequence. An antisense or sense molecule may also be directed to terminal or internal portions of the gene encoding a polypeptide having a F3'5'H activity or to combinations of the above such that the
- 15 expression of the gene is reduced or eliminated.

- With respect to this aspect of the invention, there is provided an oligonucleotide of 5-50 nucleotides such as 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48,
- 20 49, 50, 51, 52, 53, 54, 55 having substantial similarity to a part or region of a molecule with a nucleotide sequence set forth in SEQ ID NO:9 or SEQ ID NO:11 or SEQ ID NO:13 or SEQ ID NO:15 or SEQ ID NO:17 or SEQ ID NO:31 or SEQ ID NO:26. By substantial similarity or complementarity in this context is meant a hybridizable similarity under low, alternatively and preferably medium and alternatively and most preferably high stringency
- 25 conditions specific for oligonucleotide hybridization (Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2<sup>nd</sup> edition, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, USA, 1989). Such an oligonucleotide is useful, for example, in screening for F3'5'H genetic sequences from various sources or for monitoring an introduced genetic sequence in a transgenic plant. The preferred oligonucleotide is directed to a conserved F3'5'H
- 30 genetic sequence or a sequence conserved within a plant genus, plant species and/or plant variety.

- 35 -

In one aspect of the present invention, the oligonucleotide corresponds to the 5' or the 3' end of the F3'5'H genetic sequences. For convenience, the 5' end is considered herein to define a region substantially between the start codon of the structural gene to a centre portion of the gene, and the 3' end is considered herein to define a region substantially between the centre portion of the gene and the terminating codon of the structural gene. It is clear, therefore, that oligonucleotides or probes may hybridize to the 5' end or the 3' end or to a region common to both the 5' and the 3' ends. The present invention extends to all such probes.

10

In one embodiment, the nucleic acid sequence encoding a F3'5'H or various functional derivatives thereof is used to reduce the level of an endogenous a F3'5'H (e.g. *via* co-suppression or antisense-mediated suppression) or other post-transcriptional gene silencing (PTGS) processes including RNAi or alternatively the nucleic acid sequence encoding this enzyme or various derivatives or parts thereof is used in the sense or antisense orientation to reduce the level of a F3'5'H. The use of sense strands, double or partially single stranded such as constructs with hairpin loops is particularly useful in inducing a PTGS response. In a further alternative, ribozymes, minizymes or DNazymes could be used to inactivate target nucleic acid sequences.

20

Still a further embodiment encompasses post-transcriptional inhibition to reduce translation into polypeptide material. Still yet another embodiment involves specifically inducing or removing methylation.

25

Reference herein to the altering of a F3'5'H activity relates to an elevation or reduction in activity of up to 30% or more preferably of 30-50%, or even more preferably 50-75% or still more preferably 75% or greater above or below the normal endogenous or existing levels of activity. Such elevation or reduction may be referred to as modulation of a F3'5'H enzyme activity. Generally, modulation is at the level of transcription or translation of

30

F3'5'H genetic sequences.

- 36 -

The nucleic acids of the present invention may be a ribonucleic acid or deoxyribonucleic acids, single or double stranded and linear or covalently closed circular molecules. Preferably, the nucleic acid molecule is cDNA. The present invention also extends to other nucleic acid molecules which hybridize under low, preferably under medium and most preferably under high stringency conditions with the nucleic acid molecules of the present invention and in particular to the sequence of nucleotides set forth in SEQ ID NO:9 or SEQ ID NO:11 or SEQ ID NO:13 or SEQ ID NO:15 or SEQ ID NO:17 or SEQ ID NO:31 or SEQ ID NO:26 or a part or region thereof. In its most preferred embodiment, the present invention extends to a nucleic acid molecule having a nucleotide sequence set forth in SEQ ID NO:9 or SEQ ID NO:11 or SEQ ID NO:13 or SEQ ID NO:15 or SEQ ID NO:17 or SEQ ID NO:31 or SEQ ID NO:26 or to a molecule having at least 40%, more preferably at least 45%, even more preferably at least 55%, still more preferably at least 65%-70%, and yet even more preferably greater than 85% similarity at the level of nucleotide or amino acid sequence to at least one or more regions of the sequence set forth in SEQ ID NO:9 or SEQ ID NO:11 or SEQ ID NO:13 or SEQ ID NO:15 or SEQ ID NO:17 or SEQ ID NO:31 or SEQ ID NO:26 and wherein the nucleic acid encodes or is complementary to a sequence which encodes an enzyme having a F3'5'H activity. It should be noted, however, that nucleotide or amino acid sequences may have similarities below the above given percentages and yet still encode a F3'5'H activity and such molecules may still be considered in the scope of the present invention where they have regions of sequence conservation. The present invention further extends to nucleic acid molecules in the form of oligonucleotide primers or probes capable of hybridizing to a portion of the nucleic acid molecules contemplated above, and in particular those set forth in SEQ ID NO:9 or SEQ ID NO:11 or SEQ ID NO:13 or SEQ ID NO:15 or SEQ ID NO:17 or SEQ ID NO:31 or SEQ ID NO:26, under low, preferably under medium and most preferably under high stringency conditions. Preferably the portion corresponds to the 5' or the 3' end of the gene. For convenience the 5' end is considered herein to define a region substantially between the start codon of the structural genetic sequence to a centre portion of the gene, and the 3' end is considered herein to define a region substantially between the centre portion of the gene and the terminating codon of the structural genetic sequence. It is clear, therefore, that

- 37 -

oligonucleotides or probes may hybridize to the 5' end or the 3' end or to a region common to both the 5' and the 3' ends. The present invention extends to all such probes.

The term gene is used in its broadest sense and includes cDNA corresponding to the exons of a gene. Accordingly, reference herein to a gene is to be taken to include:-

- (i) a classical genomic gene consisting of transcriptional and/or translational regulatory sequences and/or a coding region and/or non-translated sequences (i.e. introns, 5'- and 3'- untranslated sequences); or
- (ii) mRNA or cDNA corresponding to the coding regions (i.e. exons) and 5'- and 3'- untranslated sequences of the gene.

The term gene is also used to describe synthetic or fusion molecules encoding all or part of an expression product. In particular embodiments, the term nucleic acid molecule and gene may be used interchangeably.

The nucleic acid or its complementary form may encode the full-length enzyme or a part or derivative thereof. By "derivative" is meant any single or multiple amino acid substitutions, deletions, and/or additions relative to the naturally occurring enzyme and which retains a F3'5'H activity. In this regard, the nucleic acid includes the naturally occurring nucleotide sequence encoding a F3'5'H or may contain single or multiple nucleotide substitutions, deletions and/or additions to said naturally occurring sequence. The nucleic acid of the present invention or its complementary form may also encode a "part" of the F3'5'H, whether active or inactive, and such a nucleic acid molecule may be useful as an oligonucleotide probe, primer for polymerase chain reactions or in various mutagenic techniques, or for the generation of antisense molecules.

Reference herein to a "part" of a nucleic acid molecule, nucleotide sequence or amino acid sequence, preferably relates to a molecule which contains at least about 10 contiguous nucleotides or five contiguous amino acids, as appropriate.

- 38 -

Amino acid insertional derivatives of the F3'5'H of the present invention include amino and/or carboxyl terminal fusions as well as intra-sequence insertions of single or multiple amino acids. Insertional amino acid sequence variants are those in which one or more amino acid residues are introduced into a predetermined site in the protein although random insertion is also possible with suitable screening of the resulting product. Deletional variants are characterized by the removal of one or more amino acids from the sequence. Substitutional amino acid variants are those in which at least one residue in the sequence has been removed and a different residue inserted in its place. Typical substitutions are those made in accordance with Table 3.

**TABLE 3** Suitable residues for amino acid substitutions

Original residue	Exemplary substitutions
Ala	Ser
Arg	Lys
Asn	Gln; His
Asp	Glu
Cys	Ser
Gln	Asn; Glu
Glu	Asp
Gly	Pro
His	Asn; Gln
Ile	Leu; Val
Leu	Ile; Val
Lys	Arg; Gln; Glu
Met	Leu; Ile; Val
Phe	Met; Leu; Tyr
Ser	Thr
Thr	Ser

- 39 -

Original residue	Exemplary substitutions
Trp	Tyr
Tyr	Trp; Phe
Val	Ile; Leu; Met

Where the F3'5'H is derivatized by amino acid substitution, the amino acids are generally replaced by other amino acids having like properties, such as hydrophobicity, hydrophilicity, electronegativity, bulky side chains and the like. Amino acid substitutions are typically of single residues. Amino acid insertions will usually be in the order of about 1-10 amino acid residues and deletions will range from about 1-20 residues. Preferably, deletions or insertions are made in adjacent pairs, i.e. a deletion of two residues or insertion of two residues.

The amino acid variants referred to above may readily be made using peptide synthetic techniques well known in the art, such as solid phase peptide synthesis (Merrifield, *J. Am. Chem. Soc.* 85: 2149, 1964) and the like, or by recombinant DNA manipulations. Techniques for making substitution mutations at predetermined sites in DNA having known or partially known sequence are well known and include, for example, M13 mutagenesis. The manipulation of DNA sequence to produce variant proteins which manifest as substitutional, insertional or deletional variants are conveniently described, for example, in Sambrook *et al.* (1989, *supra*).

Other examples of recombinant or synthetic mutants and derivatives of the F3'5'H enzyme of the present invention include single or multiple substitutions, deletions and/or additions of any molecule associated with the enzyme such as carbohydrates, lipids and/or proteins or polypeptides.

The terms "analogs" and "derivatives" also extend to any functional chemical equivalent of a F3'5'H and also to any amino acid derivative described above. For convenience, reference to F3'5'H herein includes reference to any functional mutant, derivative, part, fragment, homolog or analog thereof.

- 40 -

The present invention is exemplified using nucleic acid sequences derived from pansy, salvia, sollya or lavender or kennedia since this represents the most convenient and preferred source of material to date. However, one skilled in the art will immediately appreciate that similar sequences can be isolated from any number of sources such as other plants or certain microorganisms. All such nucleic acid sequences encoding directly or indirectly a F3'5'H are encompassed by the present invention regardless of their source. Examples of other suitable sources of genes encoding F3'5'H include, but are not limited to *Vitis spp.*, *Babiana stricta*, *Pinus spp.*, *Picea spp.*, *Larix spp.*, *Phaseolus spp.*, *Vaccinium spp.*, *Cyclamen spp.*, *Iris spp.*, *Pelargonium spp.*, *Liparidae*, *Geranium spp.*, *Pisum spp.*, *Lathyrus spp.*, *Clitoria spp.*, *Catharanthus spp.*, *Malva spp.*, *Mucuna spp.*, *Vicia spp.*, *Saintpaulia spp.*, *Lagerstroemia spp.*, *bouchina spp.*, *Plumbago spp.*, *Hypocalyptus spp.*, *Rhododendron spp.*, *Linum spp.*, *Macroptilium spp.*, *Hibiscus spp.*, *Hydrangea spp.*, *Cymbidium spp.*, *Millettia spp.*, *Hedysarum spp.*, *Lespedeza spp.*, *Asparagus spp.*, *Antigonon spp.*, *Freesia spp.*, *Brunella spp.*, *Clarkia spp.*, etc.

In accordance with the present invention, a nucleic acid sequence encoding a F3'5'H may be introduced into and expressed in a transgenic plant in either orientation thereby providing a means either to convert suitable substrates, if synthesized in the plant cell, ultimately into DHM, or alternatively to inhibit such conversion of metabolites by reducing or eliminating endogenous or existing F3'5'H activity. The production of these 3', 5'-hydroxylated substrates will subsequently be converted to delphinidin-based pigments that will modify petal color and may contribute to the production of a bluer color. Expression of the nucleic acid sequence in the plant may be constitutive, inducible or developmental and may also be tissue-specific. The word "expression" is used in its broadest sense to include production of RNA or of both RNA and protein. It also extends to partial expression of a nucleic acid molecule.

According to this aspect of the present invention, there is provided a method for producing a transgenic flowering plant capable of synthesizing a F3'5'H, said method comprising stably transforming a cell of a suitable plant with a nucleic acid sequence which comprises a sequence of nucleotides encoding said F3'5'H under conditions permitting the eventual



- 41 -

expression of said nucleic acid sequence, regenerating a transgenic plant from the cell and growing said transgenic plant for a time and under conditions sufficient to permit the expression of the nucleic acid sequence. The transgenic plant may thereby produce non-indigenous F3'5'H at elevated levels relative to the amount expressed in a comparable non-transgenic plant.

Another aspect of the present invention contemplates a method for producing a transgenic plant with reduced indigenous or existing flavonoid 3', 5'-hydroxylase activity, said method comprising stably transforming a cell of a suitable plant with a nucleic acid molecule which comprises a sequence of nucleotides encoding or complementary to a sequence encoding a F3'5'H activity, regenerating a transgenic plant from the cell and where necessary growing said transgenic plant under conditions sufficient to permit the expression of the nucleic acid.

Yet another aspect of the present invention contemplates a method for producing a genetically modified plant with reduced indigenous or existing F3'5'H activity, said method comprising altering the *F3'5'H* gene through modification of the indigenous sequences via homologous recombination from an appropriately altered *F3'5'H* gene or derivative or part thereof introduced into the plant cell, and regenerating the genetically modified plant from the cell.

As used herein an "indigenous" enzyme is one, which is native to or naturally expressed in a particular cell. A "non-indigenous" enzyme is an enzyme not native to the cell but expressed through the introduction of genetic material into a plant cell, for example, through a transgene. An "endogenous" enzyme is an enzyme produced by a cell but which may or may not be indigenous to that cell.

In a preferred embodiment, the present invention contemplates a method for producing a transgenic flowering plant exhibiting altered floral or inflorescence properties, said method comprising stably transforming a cell of a suitable plant with a nucleic acid sequence of the present invention, regenerating a transgenic plant from the cell and growing said transgenic

- 42 -

plant for a time and under conditions sufficient to permit the expression of the nucleic acid sequence.

Alternatively, said method may comprise stably transforming a cell of a suitable plant with  
5 a nucleic acid sequence of the present invention or its complementary sequence, regenerating a transgenic plant from the cell and growing said transgenic plant for a time and under conditions sufficient to alter the level of activity of the indigenous or existing F3'5'H. Preferably the altered level would be less than the indigenous or existing level of F3'5'H activity in a comparable non-transgenic plant. Without wishing to limit the present  
10 invention, one theory of mode of action is that reduction of the indigenous F3'5'H activity requires the expression of the introduced nucleic acid sequence or its complementary sequence. However, expression of the introduced genetic sequence or its complement may not be required to achieve the desired effect: namely, a flowering plant exhibiting altered floral or inflorescence properties.

15 In a related embodiment, the present invention contemplates a method for producing a flowering plant exhibiting altered floral or inflorescence properties, said method comprising alteration of the flavonoid 3', 5'-hydroxylase gene through modification of the indigenous sequences via homologous recombination from an appropriately altered F3'5'H  
20 gene or derivative or part thereof introduced into the plant cell, and regenerating the genetically modified plant from the cell.

Preferably, the altered floral or inflorescence includes the production of different shades of blue or purple or red flowers or other colors, depending on the genotype and physiological  
25 conditions of the recipient plant.

Accordingly, the present invention extends to a method for producing a transgenic plant capable of expressing a recombinant gene encoding a F3'5'H or part thereof or which carries a nucleic acid sequence which is substantially complementary to all or a part of a  
30 mRNA molecule encoding the F3'5'H, said method comprising stably transforming a cell of a suitable plant with the isolated nucleic acid molecule comprising a sequence of

- 43 -

nucleotides encoding, or complementary to a sequence encoding, a F3'5'H, where necessary under conditions permitting the eventual expression of said isolated nucleic acid molecule, and regenerating a transgenic plant from the cell. By suitable plant is meant a plant capable of producing DHK and possessing the appropriate physiological properties  
5 required for the development of the color desired.

One skilled in the art will immediately recognise the variations applicable to the methods of the present invention, such as increasing or decreasing the expression of the enzyme naturally present in a target plant leading to differing shades of colors such as different  
10 shades of blue, purple or red.

The present invention, therefore, extends to all transgenic plants or parts or cells therefrom of transgenic plants or progeny of the transgenic plants containing all or part of the nucleic acid sequences of the present invention, or antisense forms thereof and/or any homologs or related forms thereof and, in particular, those transgenic plants which exhibit altered floral  
15 or inflorescence properties. The transgenic plants may contain an introduced nucleic acid molecule comprising a nucleotide sequence encoding or complementary to a sequence encoding a F3'5'H. Generally, the nucleic acid would be stably introduced into the plant genome, although the present invention also extends to the introduction of a F3'5'H  
20 nucleotide sequence within an autonomously-replicating nucleic acid sequence such as a DNA or RNA virus capable of replicating within the plant cell. The invention also extends to seeds from such transgenic plants. Such seeds, especially if colored, are useful as proprietary tags for plants. Any and all methods for introducing genetic material into plant cells including but not limited to *Agrobacterium*-mediated transformation, biolistic particle  
25 bombardment etc. are encompassed by the present invention.

Another aspect of the present invention contemplates the use of the extracts from transgenic plants or plant parts or cells therefrom of transgenic plants or progeny of the transgenic plants containing all or part of the nucleic acid sequences of the present  
30 invention and, in particular, the extracts from those transgenic plants when used as a flavouring or food additive or health product or beverage or juice or coloring.

- 44 -

Plant parts contemplated by the present invention includes, but is not limited to flowers, fruits, vegetables, nuts, roots, stems, leaves or seeds.

- 5 The extracts of the present invention may be derived from the plants or plant part or cells therefrom in a number of different ways including but not limited to chemical extraction or heat extraction or filtration or squeezing or pulverization.

- 10 The plant, plant part or cells therefrom or extract can be utilized in any number of different ways such as for the production of a flavouring (e.g. a food essence), a food additive (e.g. a stabilizer, a colorant) a health product (e.g. an antioxidant, a tablet) a beverage (e.g. wine, spirit, tea) or a juice (e.g. fruit juice) or coloring (e.g. food coloring, fabric coloring, dye, paint, tint).

- 15 A further aspect of the present invention is directed to recombinant forms of F3'5'H. The recombinant forms of the enzyme will provide a source of material for research, for example, more active enzymes and may be useful in developing *in vitro* systems for production of colored compounds.

- 20 Still a further aspect of the present invention contemplates the use of the genetic sequences described herein in the manufacture of a genetic construct capable of expressing a F3'5'H or down-regulating an indigenous F3'5'H enzyme in a plant.

- 25 The term genetic construct has been used interchangeably throughout the specification and aligns with the terms "fusion molecule", "recombinant molecule", "recombinant nucleotide sequence". A genetic construct may include a single nucleic acid molecule comprising a nucleotide sequence encoding a single protein or may contain multiple open reading frames encoding 2 or more proteins. It may also contain a promoter operably linked to 1 or more of the open reading frames.

30

- 45 -

Another aspect of the present invention is directed to a prokaryotic or eukaryotic organism carrying a genetic sequence encoding a F3'5'H extrachromasomally in plasmid form.

5 The present invention further extends to a recombinant polypeptide comprising a sequence of amino acids substantially as set forth in SEQ ID NO:10 or SEQ ID NO:12 or SEQ ID NO:14 or SEQ ID NO:16 or SEQ ID NO:18 or SEQ ID NO:32 or SEQ ID NO:27 or an amino acid sequence having at least about 50% similarity to SEQ ID NO:10 or SEQ ID NO:12 or SEQ ID NO:14 or SEQ ID NO:16 or SEQ ID NO:18 or SEQ ID NO:32 or SEQ ID NO:27 or a derivative of said polypeptide.

10

A "recombinant polypeptide" means a polypeptide encoded by a nucleotide sequence introduced into a cell directly or indirectly by human intervention or into a parent or other relative or precursor of the cell. A recombinant polypeptide may also be made using cell-free, *in vitro* transcription systems. The term "recombinant polypeptide" includes an isolated polypeptide or when present in a cell or cell preparation. It may also be in a plant or parts of a plant regenerated from a cell which produces said polypeptide.

15

A "polypeptide" includes a peptide or protein and is encompassed by the term "enzyme".

20 The recombinant polypeptide may also be a fusion molecule comprising two or more heterologous amino acid sequences.

The present invention is further described by the following non-limiting Examples.

- 46 -

**EXAMPLE 1*****General methods***

In general, the methods followed were as described in Sambrook *et al.* (1989, *supra*) or  
 5 Sambrook and Russell, Molecular Cloning: A Laboratory Manual 3<sup>rd</sup> edition, Cold Spring  
 Harbor Laboratories, Cold Spring Harbor, NY, USA, 2001 or Plant Molecular Biology  
 Manual (2<sup>nd</sup> edition), Gelvin and Schilperoot (eds), Kluwer Academic Publisher, The  
 Netherlands, 1994 or Plant Molecular Biology Labfax, Croy (ed), Bios scientific  
 Publishers, Oxford, UK, 1993.

10

The cloning vectors pBluescript and PCR script were obtained from Stratagene, USA.  
 pCR7 2.1 was obtained from Invitrogen, USA.

***E. coli transformation***15 The *Escherichia coli* strains used were:DH5 $\alpha$ 

supE44,  $\Delta$  (lacZYA-ArgF)U169, ( $\phi$ 80lacZAM15), hsdR17( $r_k^-$ ,  $m_k^+$ ),  
 recA1, endA1, gyrA96, thi-1, relA1, deoR. (Hanahan, *J. Mol. Biol.* 166: 557, 1983)

20

XL1-Blue

supE44, hsdR17( $r_k^-$ ,  $m_k^+$ ), recA1, endA1, gyrA96, thi-1, relA1,  
 lac<sup>-</sup>, [F'proAB, lacI<sup>q</sup>, lacZAM15, Tn10(tet<sup>R</sup>)] (Bullock *et al.*, *Biotechniques* 5: 376, 1987).

25 BL21-CodonPlus-RIL strain

*ompT hsdS*(Rb- mB-) *dcm*<sup>+</sup> Tet<sup>r</sup> *gal endA Hte* [*argU ileY leuW Cam*<sup>r</sup>]

M15 *E. coli* is derived from *E. coli* K12 and has the phenotype Nal<sup>s</sup>, Str<sup>s</sup>, Rif<sup>s</sup>, Thi<sup>-</sup>, Ara<sup>+</sup>,  
 Gal<sup>+</sup>, Mtl<sup>-</sup>, F<sup>-</sup>, RecA<sup>+</sup>, Uvr<sup>+</sup>, Lon<sup>+</sup>.

30 Transformation of the *E. coli* strains was performed according to the method of Inoue *et al.*, (*Gene* 96: 23-28, 1990).

- 47 -

**Agrobacterium tumefaciens strains and transformations**

The disarmed *Agrobacterium tumefaciens* strain used was AGL0 (Lazo *et al. Bio/technology* 9: 963-967, 1991).

5

Plasmid DNA was introduced into the *Agrobacterium tumefaciens* strain AGL0 by adding 5 µg of plasmid DNA to 100 µL of competent AGL0 cells prepared by inoculating a 50 mL LB culture (Sambrook *et al.*, 1989, *supra*) and incubation for 16 hours with shaking at 28°C. The cells were then pelleted and resuspended in 0.5 mL of 85% (v/v) 100 mM CaCl<sub>2</sub>/15% (v/v) glycerol. The DNA-*Agrobacterium* mixture was frozen by incubation in liquid N<sub>2</sub> for 2 minutes and then allowed to thaw by incubation at 37°C for 5 minutes. The DNA/bacterial mix was then placed on ice for a further 10 minutes. The cells were then mixed with 1mL of LB (Sambrook *et al.*, 1989 *supra*) media and incubated with shaking for 16 hours at 28°C. Cells of *A. tumefaciens* carrying the plasmid were selected on LB agar plates containing appropriate antibiotics such as 50 µg/mL tetracycline or 100 µg/mL gentamycin. The confirmation of the plasmid in *A. tumefaciens* was done by restriction endonuclease mapping of DNA isolated from the antibiotic-resistant transformants.

10  
15**DNA ligations**

DNA ligations were carried out using the Amersham Ligation Kit or Promega Ligation Kit according to procedures recommended by the manufacturer.

20

**Isolation and purification of DNA fragments**

Fragments were generally isolated on a 1% (w/v) agarose gel and purified using the QIAEX II Gel Extraction kit (Qiagen) or Bresaclean Kit (Bresatec, Australia) following procedures recommended by the manufacturer.

25

**Repair of overhanging ends after restriction endonuclease digestion**

Overhanging 5' ends were repaired using DNA polymerase (Klenow fragment) according to standard protocols (Sambrook *et al.*, 1989 *supra*). Overhanging 3' ends were repaired using T4 DNA polymerase according to standard protocols (Sambrook *et al.*, 1989 *supra*).

30

- 48 -

**Removal of phosphoryl groups from nucleic acids**

Shrimp alkaline phosphatase (SAP) (USB) was typically used to remove phosphoryl groups from cloning vectors to prevent re-circularization according to the manufacturer's recommendations.

**Polymerase Chain Reaction (PCR)**

Unless otherwise specified, PCR conditions using plasmid DNA as template included using 2 ng of plasmid DNA, 100 ng of each primer, 2  $\mu$ L 10 mM dNTP mix, 5  $\mu$ L 10 x Taq DNA polymerase buffer, 0.5  $\mu$ L Taq DNA Polymerase in a total volume of 50  $\mu$ L. Cycling conditions comprised an initial denaturation step of 5 minutes at 94°C, followed by 35 cycles of 94°C for 20 sec, 50°C for 30 sec and 72°C for 1 minute with a final treatment at 72°C for 10 minutes before storage at 4°C.

PCRs were performed in a Perkin Elmer GeneAmp PCR System 9600.

**<sup>32</sup>P-Labeling of DNA Probes**

DNA fragments (50 to 100 ng) were radioactively labelled with 50  $\mu$ Ci of [ $\alpha$ -<sup>32</sup>P]-dCTP using a Gigaprime kit (Geneworks). Unincorporated [ $\alpha$ -<sup>32</sup>P]-dCTP was removed by chromatography on Sephadex G-50 (Fine) columns or Microbiospin P-30 Tris chromatography columns (BioRad).

**Plasmid Isolation**

Single colonies were analyzed for inserts by inoculating LB broth (Sambrook *et al.*, 1989, *supra*) with appropriate antibiotic selection (e.g. 100  $\mu$ g/mL ampicillin or 10 to 50  $\mu$ g/mL tetracycline etc.) and incubating the liquid culture at 37°C (for *E. coli*) or 29°C (for *A. tumefaciens*) for ~16 hours with shaking. Plasmid DNA was purified using the alkali-lysis procedure (Sambrook *et al.*, 1989, *supra*) or using The WizardPlus SV minipreps DNA purification system (Promega) or Qiagen Plasmid Mini Kit (Qiagen). Once the presence of an insert had been determined, larger amounts of plasmid DNA were prepared from 50 mL overnight cultures using the alkali-lysis procedure (Sambrook *et al.*, 1989, *supra*) or



- 49 -

QIAfilter Plasmid Midi kit (Qiagen) and following conditions recommended by the manufacturer.

**DNA Sequence Analysis**

- 5 DNA sequencing was performed using the PRISM (trademark) Ready Reaction Dye Primer Cycle Sequencing Kits from Applied Biosystems. The protocols supplied by the manufacturer were followed. The cycle sequencing reactions were performed using a Perkin Elmer PCR machine (GeneAmp PCR System 9600). Sequencing runs were generally performed by the Australian Genome Research Facility at The Walter and Eliza  
10 Hall Institute of Medical Research (Melbourne, Australia) or in-house on an automated 373A DNA sequencer (Applied Biosystems).

Sequences were analysed using a MacVector™ application (version 6.5.3) (Oxford Molecular Ltd., Oxford, England).

15

- Homology searches against Genbank, SWISS-PROT and EMBL databases were performed using the FASTA and TFASTA programs (Pearson and Lipman, *Proc. Natl. Acad. Sci. USA* 85(8): 2444-2448, 1988) or BLAST programs (Altschul *et al.*, *J. Mol. Biol.* 215(3): 403-410, 1990). Percentage sequence similarities were obtained using LALIGN  
20 program (Huang and Miller, *Adv. Appl. Math.* 12: 373-381, 1991) or ClustalW program (Thompson *et al.*, *Nucleic Acids Research* 22: 4673-4680, 1994) within the MacVector™ application (Oxford Molecular Ltd., England) using default settings.

- Multiple sequence alignments were produced using ClustalW (Thompson *et al.*, 1994,  
25 *supra*) using default settings.

- 50 -

**EXAMPLE 2*****Plant transformations******Petunia hybrida* transformations (Sw63 x Skr4)**

- 5 As described in Holton *et al.* (1993a, *supra*) by any other method well known in the art.

***Rosa hybrida* transformations**

- As described in U.S. Patent Application No. 542,841 (PCT/US91/04412) or Robinson and Firoozabady (*Scientia Horticulturae*, 55: 83-99, 1993), Rout *et al.* (*Scientia Horticulturae*,  
10 81: 201-238, 1999) or Marchant *et al.* (*Molecular Breeding* 4: 187-194, 1998) or by any other method well known in the art.

Cuttings of *Rosa hybrida* were generally obtained from Van Wyk and Son Flower Supply, Victoria.

15

***Dianthus caryophyllus* transformations**

- International Patent Application No. PCT/US92/02612 (carnation transformation). As described in International Patent Application No. PCT/AU96/00296 (Violet carnation), Lu  
20 *et al.* (*Bio/Technology* 9: 864-868, 1991), Robinson and Firoozabady (1993, *supra*) or by any other method well known in the art.

Cuttings of *Dianthus caryophyllus* cv. Kortina Chanel or Monte Lisa were obtained from Van Wyk and Son Flower Supply, Victoria.

25

- 51 -

**EXAMPLE 3*****Transgenic Analysis*****Color coding**

- 5 The Royal Horticultural Society's Color Chart (Kew, UK) was used to provide a description of observed color. They provide an alternative means by which to describe the color phenotypes observed. The designated numbers, however, should be taken only as a guide to the perceived colors and should not be regarded as limiting the possible colors which may be obtained.

10

**Chromatographic analysis**

Thin Layer Chromatography (TLC) and High Performance Liquid Chromatography (HPLC) analysis was performed generally as described in Brugliera *et al.* (*Plant J.* 5, 81-92, 1994).

15

**Extraction of anthocyanidins**

- Prior to HPLC analysis, the anthocyanin and flavonol molecules present in petal and stamen extracts were acid hydrolysed to remove glycosyl moieties from the anthocyanidin or flavonol core. Anthocyanidin and flavonol standards were used to help identify the compounds present in the floral extracts.

20

- Anthocyanidins in the reaction mixture were analysed by HPLC *via* gradient elution using gradient conditions of 50%B to 60%B over 10 minutes, then 60% B for 10 minutes and finally 60% B to 100% B over 5 minutes where solvent A consisted of TFA: H<sub>2</sub>O (5:995) and solvent B consisted of acetonitrile: TFA: H<sub>2</sub>O (500:5:495). An Asahi Pac ODP-50 cartridge column (250 mm x 4.6 mm ID) was used for the reversed phase chromatographic separations. The flow rate was 1 mL/min and the temperature was 40°C. The detection of the anthocyanidin compounds was carried out using a Shimadzu SPD-M6A three dimensional detector at 400-650 nm.

30

- 52 -

The anthocyanidin peaks were identified by reference to known standards, viz delphinidin or delphinidin-based molecules, petunidin, malvidin, cyanidin and peonidin

**Stages of flower development**

5

***Petunia***

*Petunia hybrida* cv. Skr4 x Sw63 flowers were harvested at developmental stages defined as follows:

- 10    Stage 1:        Unpigmented, closed bud.  
      Stage 2:        Pigmented, closed bud.  
      Stage 3:        Pigmented bud with emerging corolla  
      Stage 4:        Pigmented, opened flower with anther intact (pre-dehiscence)  
      Stage 5:        Fully opened flower with all anthers dehisced.

15

For TLC or HPLC analysis, petals were collected from stage 4 flowers at the stage of maximum pigment accumulation.

- For Northern blot analysis, petals were collected from stages 2 to 3 flowers at the stage of  
20    maximal expression of flavonoid pathway genes.

***Carnation***

*Dianthus caryophyllus* flowers were harvested at developmental stages defined as follows:

- 25    Stage 1:        Closed bud, petals not visible.  
      Stage 2:        Flower buds opening: tips of petals visible.  
      Stage 3:        Tips of nearly all petals exposed. "Paint-brush stage".  
      Stage 4:        Outer petals at 45° angle to stem.  
      Stage 5:        Flower fully open.

30

- 53 -

For TLC or HPLC analysis, petals were collected from stage 4 flowers at the stage of maximum pigment accumulation.

For Northern blot analysis, petals were collected from stage 3 flowers at the stage of maximal expression of flavonoid pathway genes.

#### *Rose*

Stages of *Rosa hybrida* flower development were defined as follows:

- 10 Stage 1: Unpigmented, tightly closed bud.
- Stage 2: Pigmented, tightly closed bud
- Stage 3: Pigmented, closed bud; sepals just beginning to open.
- Stage 4: Flower bud beginning to open; petals heavily pigmented; sepals have separated.
- 15 Stage 5: Sepals completely unfolded; some curling. Petals are heavily pigmented and unfolding.

For TLC or HPLC analysis, petals were collected from stage 4 flowers at the stage of maximum pigment accumulation.

20

For Northern blot analysis, petals were collected from stage 3 to 4 flowers at the stage of maximal expression of flavonoid pathway genes (Tanaka *et al.*, *Plant Cell Physiol.*, 36(6): 1023-1031, 1995).

#### 25 *Anthocyanin/flavonol measurements by spectrophotometric measurements*

Approximately 200mg of fresh petal tissue was added to 2 mL of methanol/1% (v/v) HCl and incubated for ~16 hours at 4°C. A 1 in 20 dilution (e.g. 50 µL made to 1000 µL) was then made and the absorbance at 350 nm and 530 nm was recorded.

30 The approximate flavonols and anthocyanin amounts (nmoles/gram) were then calculated according to the following formulae:

*Anthocyanin content*

$$\frac{(A_{530} / 34,000) \times \text{volume of extraction buffer (mL)} \times \text{dilution factor} \times 10^6}{\text{mass of petal tissue (grams)}}$$

5

*Flavonol content*

$$\frac{(A_{350} / 14,300) \times \text{volume of extraction buffer (mL)} \times \text{dilution factor} \times 10^6}{\text{mass of petal tissue (grams)}}$$

10 *Northern/RNA blot analysis*

Transcription of a transferred gene was monitored by isolating RNA and estimating the quantity and size of the expected transcript. Northern blot analysis was used to monitor the steady-state level of particular transcripts in petals. A transcript was determined to be intact or full-length based on the estimated size expected from the gene used. In general

15 when cDNAs were used as coding sequences the size of the transcript expected would be the size of the cDNA plus any 5' untranslated component of the fused promoter fragment plus any 3' untranslated sequence from the fused terminator fragment. In some cases where a cDNA region contained a putative polyadenylation site and the terminator region contained a putative polyadenylation site, 2 transcripts would be detected. One would be of

20 a size consistent with polyadenylation occurring just downstream from the polyadenylation site within the cDNA sequence. The second transcript would be larger and consistent with the transcript being polyadenylated after the polyadenylation site within the terminator fragment.

25 Total RNA was isolated from petals or leaves using a Plant RNAeasy kit (QIAGEN) following procedures recommended by the manufacturer. For rose samples 1% (w/v) PVP was added to the extraction buffer.

- 55 -

RNA samples (5 µg) were electrophoresed through 2.2 M formaldehyde/1.2% w/v agarose gels using running buffer containing 40 mM morpholinopropanesulphonic acid (pH 7.0), 5 mM sodium acetate, 0.1 mM EDTA (pH 8.0). The RNA was stained with ethidium bromide and visualised under UV-light. The ribosomal RNA was generally used as a guide  
5 in confirming that the RNA had not been degraded by intra- or extra- cellular ribonucleases. The RNA was transferred to Hybond-N membrane filters (Amersham) and treated as described by the manufacturer.

Control samples were included on RNA gels as a measure of the integrity of the  
10 radiolabelled probe and as guides to expected transcript sizes. Controls for *petHf1* and *petHf2* genes included RNA isolated from petunia OGB petals (stages 3 to 4) or from flowers of transgenic carnations shown previously to accumulate *petHf1* transcripts. Controls for other *F3'5'H* genes generally included RNA isolated from petals of the same species from which the *F3'5'H* sequence had been isolated.

15 RNA blots were probed with <sup>32</sup>P-labelled fragments. Prehybridization (1 hour at 42°C) and hybridization (16 hours at 42°C) of the membrane filters were carried out in 50% v/v formamide, 1 M NaCl, 1% w/v SDS, 10% w/v dextran sulphate. The membrane filters were generally washed in 2 x SSC, 1% w/v SDS at 65°C for between 1 to 2 hours and then  
20 0.2 x SSC, 1% w/v SDS at 65°C for between 0.5 to 1 hour. Membrane filters were generally exposed to Kodak XAR film with an intensifying screen at -70°C for 16 to 72 hours.

#### EXAMPLE 4

##### 25 *Introduction of chimeric petunia F3'5'H genes into rose*

As described in the introduction, the pattern of hydroxylation of the B-ring of the anthocyanidin molecule plays a key role in determining petal color. The production of the dihydroflavonol DHM, leads to the production of the purple/blue delphinidin-based  
30 pigments in plants such as petunia. The absence of the *F3'5'H* activity has been correlated

- 56 -

with the absence of blue flowers in many plant species such as *Rosa*, *Gerbera*, *Antirrhinum*, *Dianthus* and *Dendranthema*.

Based on success in producing delphinidin-based pigments in a mutant petunia line (Holton *et al.*, 1993a, *supra* and International Patent Application No. PCT/AU92/00334), in tobacco flowers (International Patent Application No. PCT/AU92/00334) and in carnation flowers (International Patent Application No. PCT/AU96/00296), similar chimeric petunia *F3'5'H* genes were also introduced into roses in order to produce novel delphinidin-based pigments and modify flower color.

**Preparation of chimeric petunia *F3'5'H* gene constructs**

A summary of promoter, terminator and coding fragments used in the preparation of constructs and the respective abbreviations is listed in Table 4.

**TABLE 4** Abbreviations used in construct preparations

ABBREVIATION	DESCRIPTION
<i>AmCHS 5'</i>	1.2 kb promoter fragment from the <i>Antirrhinum majus</i> chalcone synthase ( <i>CHS</i> ) gene (Sommer and Saedler, <i>Mol Gen. Gen.</i> , 202: 429-434, 1986)
<i>CaMV 35S</i>	~0.2 kb incorporating <i>Bgl</i> II fragment containing the promoter region from the Cauliflower Mosaic Virus 35S ( <i>CaMV 35S</i> ) gene - (Franck <i>et al.</i> , <i>Cell</i> 21: 285-294, 1980, Guilley <i>et al.</i> , <i>Cell</i> , 30: 763-773, 1982)
<i>35S 5'</i>	promoter fragment from <i>CaMV 35S</i> gene (Franck <i>et al.</i> , 1980, <i>supra</i> ) with an ~60bp 5' untranslated leader sequence from the petunia chlorophyll a/b binding protein gene ( <i>Cab 22</i> gene) (Harpster <i>et al.</i> , <i>MGG</i> , 212: 182-190, 1988)
<i>chrysCHS 5'</i>	promoter region from a <i>CHS</i> gene from chrysanthemum (SEQ ID NO: 30)



ABBREVIATION	DESCRIPTION
<i>eCaMV 35S</i>	enhanced <i>CaMV 35S</i> promoter as described in Mitsuhashi <i>et al.</i> , <i>Plant Cell Physiol.</i> 37: 49-59, 1996
<i>GUS</i>	$\beta$ -glucuronidase ( <i>GUS</i> ) coding sequence (Jefferson, <i>et al.</i> , <i>EMBO J.</i> 6: 3901-3907, 1987)
<i>Mac</i>	Hybrid promoter consisting of the promoter from the mannopine synthase ( <i>mas</i> ) gene and a <i>CaMV 35S</i> enhancer region (Comai <i>et al.</i> , <i>Plant Mol. Biol.</i> 15: 373-381, 1990)
<i>mas/35S</i>	Hybrid promoter consisting of a promoter region from <i>CaMV 35S</i> gene with enhancer elements from a promoter fragment of mannopine synthase ( <i>mas</i> ) gene of <i>Agrobacterium tumefaciens</i> (Janssen and Gardner, <i>Plant Molecular Biology</i> , 14: 61-72, 1989)
<i>mas 5'</i>	Promoter region from the <i>mas</i> of <i>A. tumefaciens</i>
<i>mas 3'</i>	Terminator region from the <i>mas</i> gene of <i>A. tumefaciens</i>
<i>nos 5'</i>	Promoter region from the nopaline synthase ( <i>nos</i> ) gene of <i>A. tumefaciens</i> (Depicker <i>et al.</i> , <i>J Mol. and Appl. Genetics</i> 1: 561-573, 1982)
<i>nos 3'</i>	Terminator region from the <i>nos</i> gene of <i>A. tumefaciens</i> (Depicker <i>et al.</i> , 1982, <i>supra</i> )
<i>nptII</i>	Kanamycin-resistance gene (encodes neomycin phosphotransferase which deactivates aminoglycoside antibiotics such as kanamycin, neomycin and G418)
<i>ocs 3'</i>	~1.6kb terminator fragment from octopine synthase gene of <i>A. tumefaciens</i> (described in Janssen and Gardner, 1989, <i>supra</i> )
<i>petD8 5'</i>	~3.2kb promoter region from a phospholipid transfer protein gene ( <i>D8</i> ) of <i>Petunia hybrida</i> (Holton, Isolation and characterization of petal specific genes from <i>Petunia hybrida</i> . PhD thesis, University of Melbourne, Australia, 1992) (SEQ ID NO: 24)
<i>petD8 3'</i>	~0.7kb terminator region from a phospholipid transfer protein gene ( <i>D8</i> ) of <i>Petunia hybrida</i> cv. OGB (Holton, 1992, <i>supra</i> )

ABBREVIATION	DESCRIPTION
<i>long petFLS 5'</i>	~4.0kb fragment containing the promoter region from a flavonol synthase ( <i>FLS</i> ) gene of <i>P. hybrida</i>
<i>short petFLS 5'</i>	~2.2kb fragment containing the promoter region from <i>FLS</i> gene of <i>P. hybrida</i>
<i>petFLS 3'</i>	~0.95kb fragment containing the terminator region from <i>FLS</i> gene of <i>P. hybrida</i>
<i>petHf1</i>	Petunia <i>F3'5'H Hf1</i> cDNA clone (Holton <i>et al.</i> , 1993a, <i>supra</i> ) (SEQ ID NO: 1)
<i>petHf2</i>	Petunia <i>F3'5'H Hf2</i> cDNA clone (Holton <i>et al.</i> , 1993a, <i>supra</i> ) (SEQ ID NO: 3)
<i>petRT 5'</i>	Promoter region of an anthocyanidin-3- glucoside rhamnosyltransferase ( <i>3RT</i> ) gene from <i>P. hybrida</i> (Brugliera, Characterization of floral specific genes isolated from <i>Petunia hybrida</i> . RMIT, Australia. PhD thesis, 1994)
<i>petRT 3'</i>	Terminator region of a <i>3RT</i> gene from <i>P. hybrida</i> (Brugliera, 1994, <i>supra</i> )
<i>RoseCHS 5'</i>	~2.8kb fragment containing the promoter region from a <i>CHS</i> gene of <i>Rosa hybrida</i> (SEQ ID: 5)
<i>SuRB</i>	Chlorsulfuron-resistance gene (encodes Acetolactate Synthase) with its own terminator from <i>Nicotiana tabacum</i> (Lee <i>et al.</i> , <i>EMBO J.</i> 7: 1241-1248, 1988)

In order to produce delphinidin or delphinidin-based molecules in rose petals, a number of binary vector constructs were prepared utilising the petunia *F3'5'H* cDNA fragments and various promoter and terminator fragments. The chimaeric petunia *F3'5'H* genes had proved successful in carnation and petunia leading to detectable intact *F3'5'H* transcripts (as detected by Northern blot analysis) and to the production of delphinidin or delphinidin-based molecules pigments. Table 5 summarises the list of binary vector constructs containing petunia *F3'5'H* cDNA fragments.

- 59 -

**TABLE 5** Summary of chimaeric petunia *F3'5'H* gene expression cassettes contained in binary vector constructs used in the transformation of roses (see Table 4 for an explanation of abbreviations).

PLASMID	<i>F3'5'H</i> GENE	SELECTABLE MARKER GENE
pCGP1452	<i>AmCHS 5': petHf1: petD8 3'</i>	<i>35S 5': SuRB</i>
pCGP1453	<i>Mac: petHf1: mas 3'</i>	<i>35S 5': SuRB</i>
pCGP1457	<i>petD8 5': petHf1: petD8 3'</i>	<i>35S 5': SuRB</i>
pCGP1461	<i>short petFLS 5': petHf1: petFLS 3'</i>	<i>35S 5': SuRB</i>
pCGP1616	<i>petRT 5': petHf1: nos 3'</i>	<i>35S 5': SuRB</i>
pCGP1638	<i>CaMV 35S: petHf1: ocs 3'</i>	<i>35S 5': SuRB</i>
pCGP1623	<i>mas 35S: petHf1: ocs 3'</i>	<i>35S 5': SuRB</i>
pCGP1860	<i>RoseCHS 5': petHf1: nos 3'</i>	<i>35S 5': SuRB</i>
pCGP2123	<i>CaMV 35S: petHf2: ocs 3'</i>	<i>35S 5': SuRB</i>

5

**Isolation of petunia *F3'5'H* cDNA clones (*petHf1* and *petHf2*)**

The isolation and characterisation of cDNA clones of petunia *F3'5'H* (*petHf1* and *petHf2* contained in pCGP602 (Figure 2) and pCGP175 (Figure 3) respectively) (SEQ ID NO:1 and SEQ ID NO:3, respectively) have been described in International Patent Application  
10 No. PCT/AU92/00334 and Holton *et al.* (1993a, *supra*).

The plasmids pCGP601 (Figure 2), pCGP602 (Figure 2), pCGP176 (Figure 2) contain homologs of the petunia *petHf1* *F3'5'H* cDNA clone. The plasmid pCGP601 contains a petunia *F3'5'H petHf1* homolog that includes 52bp of 5' untranslated sequence. The  
15 plasmid pCGP602 contains a petunia *F3'5'H petHf1* homolog that includes 125bp of 5' untranslated sequence (SEQ ID NO:1). The plasmid pCGP176 (described in Holton *et al.*, 1993a *supra*) contains a petunia *F3'5'H petHf1* homolog that includes 27bp of 5' untranslated sequence and a further ~127bp of 3' untranslated sequence over the petunia *F3'5'H petHf1* cDNA clone in pCGP602.

20

- 60 -

**Construction of pCGP1303 (*petHf1* in *pUC19* backbone)**

The petunia *F3'5'H* cDNA clone contained in the plasmid pCGP601 (described above) (Figure 2) included 52 bp of 5' untranslated sequence and 141 bp of 3' untranslated sequence including 16 bp of the poly A tail. The plasmid pCGP601 (Figure 2) was firstly  
5 linearized by digestion with the restriction endonuclease *BspHI*. The ends were repaired and the petunia *F3'5'H petHf1* cDNA clone was released upon digestion with the restriction endonuclease *FspI*. The *BspHI* recognition sequence encompasses the putative translation initiating codon and the *FspI* recognition sequence commences 2 bp downstream from the stop codon. The 1.6 kb fragment containing the petunia *F3'5'H*  
10 *petHf1* cDNA clone was purified and ligated with repaired *EcoRI* ends of pUC19 (New England Biolabs). Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP1303 (Figure 4).

**Construction of pCGP627 (short *petHf1* in *pBluescript* backbone)**

The plasmid pCGP176 (Holton *et al.*, 1993a, *supra*) (Figure 2) was digested with the restriction endonuclease *SpeI* and *EcoRI*. The ends were then repaired and allowed to religate. The resulting plasmid was designated as pCGP627 and contained the identical cDNA clone as in pCGP176 except that the restriction endonuclease sites *PstI*, *BamHI* and  
20 *SmaI* were removed from the multi-cloning site of the pBluescript vector at the 5' end of the cDNA clone.

**The binary vector pCGP1452 (*AmCHS* 5': *petHf1*: *petD8* 3')**

The plasmid pCGP1452 (Figure 5) contains a chimaeric petunia *F3'5'H* (*petHf1*) gene under the control of a promoter fragment from the *Antirrhinum majus* chalcone synthase gene (*CHS*) (Sommer and Saedler, 1986, *supra*) with a terminator fragment from the petunia phospholipid transfer protein (PLTP) gene (*petD8* 3') (Holton, 1992, *supra*). The chimeric petunia *F3'5'H* cassette is in a tandem orientation with respect to the 35S 5': *SuRB* gene of the binary vector, pWTT2132 (DNA Plant Technologies, USA = DNAP)  
30 (Figure 6).

- 61 -

***Intermediates in the preparation of the binary pCGP1452******The binary vector pWTT2132***

The binary vector plasmid pWTT2132 (DNAP) (Figure 6) contains a chimeric gene comprised of a 35S 5' promoter sequence (Franck *et al.*, 1980, *supra*), ligated with the coding region and terminator sequence for acetolactate synthase (ALS) gene from the *SuRB* locus of tobacco (Lee *et al.*, 1988, *supra*). An ~60bp 5' untranslated leader sequence from the petunia chlorophyll a/b binding protein gene (Cab 22 gene) (Harpster *et al.*, MGG, 212: 182-190, 1988) is included between the 35S 5' promoter fragment and the *SuRB* sequence.

***Construction of pCGP725 (AmCHS 5': petHfl: petD8 3' in pBluescript)***

A chimeric petunia *F3'5'H* gene under the control *Antirrhinum majus* *CHS* (*AmCHS* 5') promoter with a petunia PLTP terminator (*petD8* 3') was constructed by cloning the 1.6kb *BclI/FspI* petunia *F3'5'H* (*petHfl*) fragment from pCGP602 (Holton *et al.*, 1993a, *supra*) (Figure 2) between a 1.2 kb *Antirrhinum majus* *CHS* gene fragment 5' to the site of translation initiation (Sommer and Saedler, 1986, *supra*) and a 0.7 kb *SmaI/XhoI* PLTP fragment (*petD8* 3') from pCGP13ΔBam (Holton, 1992, *supra*), 3' to the deduced stop codon. The resulting plasmid in a pBluescript II KS (Stratagene, USA) backbone vector was designated pCGP725 (Figure 7).

***Construction of pCGP485 and pCGP1452 (AmCHS 5': petHfl: petD8 3' binary vectors)***

The chimeric *F3'5'H* gene from pCGP725 (Figure 7) was cloned into the binary vector pCGN1547 containing an *nptII* selectable marker gene cassette (McBride and Summerfelt *Plant Molecular Biology* 14: 269-276, 1990) to create pCGP485. A 3.5 kb fragment containing the *AmCHS* 5': *petHfl*: *petD8* 3' cassette was released upon digestion of pCGP485 with the restriction endonuclease *PstI*. The overhanging ends were repaired and the purified 3.5 kb fragment was ligated with *SmaI* ends of the binary vector, pWTT2132 (DNAP). Correct insertion of the fragment in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The plasmid was designated as pCGP1452 (Figure 5).

- 62 -

**Plant transformation with pCGP1452**

The T-DNA contained in the binary vector plasmid pCGP1452 (Figure 5) was introduced into rose via *Agrobacterium*-mediated transformation.

5

**The binary vector pCGP1453 (Mac: *petHf1*: mas 3')**

The plasmid pCGP1453 (Figure 8) contains a chimeric petunia *F3'5'H* (*petHf1*) gene under the control of a *Mac* promoter (Comai *et al.*, 1990, *supra*) with a terminator fragment from the mannopine synthase gene of *Agrobacterium* (*mas 3'*). The chimeric petunia *F3'5'H* cassette is in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector, pWTT2132 (DNAP) (Figure 6).

10

A 3.9 kb fragment containing the *Mac: petHf1: mas 3'* gene was released from the plasmid pCGP628 (described in International Patent Application No. PCT/AU94/00265) upon digestion with the restriction endonuclease *Pst*I. The overhanging ends were repaired and the purified fragment was ligated with *Sma*I ends of pWTT2132 (DNAP). Correct insertion of the *Mac: petHf1: mas 3'* gene in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The plasmid was designated as pCGP1453 (Figure 8).

15

20

**Plant transformation with pCGP1453**

The T-DNA contained in the binary vector plasmid pCGP1453 (Figure 8) was introduced into rose via *Agrobacterium*-mediated transformation.

25

- 63 -

**The binary vector pCGP1457 (petD8 5': petHfl: pet D8 3')**

The plasmid pCGP1457 (Figure 9) contains a chimaeric petunia *F3'5'H* (*petHfl*) gene under the control of a promoter fragment from the petunia *PLTP* gene (*petD8 5'*) with a terminator fragment from the petunia *PLTP* gene (*petD8 3'*). The chimeric petunia *F3'5'H* cassette is in a tandem orientation with respect to the 35S 5': *SuRB* gene of the binary vector, pWTT2132 (DNAP) (Figure 6).

**Intermediates in the preparation of the binary vector pCGP1457**

**Isolation of petunia D8 genomic clone**

**10 Preparation of *P. hybrida* cv. OGB (Old Glory Blue) genomic library in  $\lambda$ 2001**

A genomic DNA library was constructed from *Petunia hybrida* cv. OGB DNA in the vector  $\lambda$ 2001 (Karn *et al.*, *Gene* 32: 217-224, 1984) using a *Sau*3A partial digestion of the genomic DNA as described in Holton, 1992 (*supra*). Screening of the OGB genomic library for the petunia D8 gene was as described in Holton, 1992. *supra*.

15

**Isolation of D8 genomic clone OGB2.6**

PCR was performed in order to find a non-mutant genomic clone representing D8. Oligo #2 (5' to 3' GTTCTCGAGGAAAGATAATAACAAT) (SEQ ID NO:6) and Oligo #4 (5' to 3' CAAGATCGTAGGACTGCATG) (SEQ ID NO:7) were used to amplify D8 gene fragments, across the intron region, using 4  $\mu$ L of phage suspension from the clones isolated from the primary screening of the OGB genomic library. The reactions were carried out in a total volume of 50  $\mu$ L containing 1 x Amplification buffer (Cetus), 0.2 mM dNTP mix, <1  $\mu$ g of template DNA, 50 pmoles of each primer and 0.25  $\mu$ L of Taq polymerase (5 units/ $\mu$ L - Cetus). The reaction mixtures were overlaid with 30  $\mu$ L of mineral oil and temperature cycled using a Gene Machine (Innovonics). The reactions were cycled 30 times using the following conditions: 94°C for 1 minute, 55°C for 50 seconds, 72°C for 2 minutes. One quarter of each PCR reaction was run on an agarose gel using TAE running buffer.

Three clones,  $\lambda$ OGB-2.4,  $\lambda$ OGB-2.5, and  $\lambda$ OGB-2.6, gave fragments of approximately 1 kb whereas the mutant clone,  $\lambda$ OGB-3.2 (described in Holton, 1992, *supra*), had produced a product of 1.25 kb. The  $\lambda$ OGB-2.6 clone was chosen for further analysis.

30

**pCGP382**

The genomic clone,  $\lambda$ OGB-2.6, contained a single 3.9 kb *Xba*I fragment that hybridized with the D8 cDNA. This *Xba*I fragment was isolated and purified and ligated with the *Xba*I ends of pBluescriptII KS- (Stratagene, USA). Restriction mapping of this clone revealed an internal *Pst*I site 350 bp from the 3' end. However, the "mutant" genomic clone in pCGP13, had an internal *Pst*I near the putative initiating "ATG" of the coding region (approximately 1.5 kb from its 3' end). The difference in the position of the *Pst*I site in both clones suggested that the  $\lambda$ OGB-2.6 *Xba*I fragment did not contain the whole genomic sequence of D8. A Southern blot was performed on *Pst*I digested  $\lambda$ OGB-2.6 DNA, and a fragment of 2.7 kb was found to hybridize with the D8 cDNA. Restriction endonuclease mapping confirmed that this fragment contained the 3' coding region and flanking sequences.

In order to obtain a fragment containing the whole D8 genomic sequence, a number of cloning steps were undertaken. The  $\lambda$ OGB-2.6 *Pst*I fragment of 2.7 kb was purified and ligated with *Pst*I ends of pBluescriptII KS- (Stratagene, USA). The resultant clone was digested with *Xba*I to remove the 350 bp *Pst*I/*Xba*I fragment. This fragment was replaced by the 3.9 kb *Xba*I fragment from  $\lambda$ OGB-2.6 to produce the plasmid pCGP382.

A 3.2 kb fragment containing the promoter region from the D8 2.6 gene in pCGP382 was released upon digestion with the restriction endonucleases *Hind*III and *Nco*I. The fragment was purified and ligated with the 4.8 kb *Nco*I/*Hind*III fragment of pJB1 (Bodeau, Molecular and genetic regulation of Bronze-2 and other maize anthocyanin genes. Dissertation, Stanford University, USA, 1994) to produce pCGP1101 containing a *petD85'*: *GUS*: *nos* 3' cassette.

A 1.6 kb petunia *F3'5'H petHf1* fragment was released from the plasmid pCGP602 (Holton *et al.*, 1993a, *supra*) (SEQ ID NO:1) (Figure 2) upon digestion with the restriction endonucleases *Bsp*HI and *Bam*HI. The fragment was purified and ligated with the 6.2 kb *Nco*I/*Bam*HI fragment of pCGP1101 to produce pCGP1102 containing a *petD8 5'*: *petHf1*: *nos* 3' expression cassette.



- 65 -

A 0.75 kb *Bam*HI *petD8* 3' fragment (Holton, 1992, *supra*) was purified from the plasmid pCGP13Δ*Bam*HI and ligated with *Bam*HI/*Bgl*II ends of pCGP1102 to produce the plasmid pCGP1107 containing a *petD8* 5': *petHf1*: *petD8* 3' expression cassette.

5

The plasmid pCGP1107 was linearised upon digestion with the restriction endonuclease *Xba*I. The overhanging ends were repaired and then the 5.3 kb fragment containing the *petD8* 5': *petHf1*: *petD8* 3' expression cassette was released upon digestion with the restriction endonuclease *Pst*I. The fragment was purified and ligated with *Sma*I/*Pst*I ends of the binary vector pWTT2132 (DNAP) (Figure 6). Correct insertion of the *petD8* 5': *petHf1*: *petD8* 3' gene in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The plasmid was designated as pCGP1457 (Figure 9).

15

**Plant transformation with pCGP1457**

The T-DNA contained in the binary vector plasmid pCGP1457 (Figure 9) was introduced into rose via *Agrobacterium*-mediated transformation.

20

**The binary vector pCGP1461 (short *petFLS* 5': *petHf1*: *pet FLS* 3')**

The plasmid pCGP1461 (Figure 10) contains a chimeric petunia *F3'5'H* (*petHf1*) gene under the control of a promoter fragment from the petunia flavonol synthase (*FLS*) gene (short *petFLS* 5') with a terminator fragment from the petunia *FLS* gene (*petFLS* 3'). The chimeric petunia *F3'5'H* gene is in a tandem orientation with respect to the 35S 5': *SuRB* gene of the binary vector, pWTT2132 (Figure 6).

25

- 66 -

***Intermediates in the preparation of the binary vector pCGP1461*****Isolation of petunia *FLS* gene*****Preparation of *P. hybrida* cv. Th7 genomic library***

A *P. hybrida* cv. Th7 genomic library was prepared according to Sambrook *et al.* (1989, 5 *supra*) using a *Sau*3A partial digestion of the genomic DNA. The partially digested DNA was cloned into EMBL-3 lambda vector (Stratagene, USA).

The Th7 genomic DNA library was screened with <sup>32</sup>P-labelled fragments of a petunia *FLS* 10 cDNA clone (Holton *et al.*, *Plant J.* 4: 1003-1010, 1993b) using high stringency conditions.

Two genomic clones (*FLS2* and *FLS3*) were chosen for further analysis and found to 15 contain sequences upstream of the putative initiating methionine of the petunia *FLS* coding region with *FLS2* containing a longer promoter region than *FLS3*.

***pCGP486***

A 6 kb fragment was released upon digestion of the genomic clone *FLS2* with the 20 restriction endonuclease *Xho*I. The fragment containing the short petunia *FLS* gene was purified and ligated with *Xho*I ends of pBluescript SK (Stratagene, USA). Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP486.

***pCGP487***

A 9 kb fragment was released upon digestion of the genomic clone *FLS3* with the 25 restriction endonuclease *Xho*I. The fragment containing the petunia *FLS* gene was purified and ligated with *Xho*I ends of pBluescript SK (Stratagene, USA). Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP487.

***pCGP717***

A 2.2 kb petunia *FLS* promoter fragment upstream from the putative translational initiation site was released from the plasmid pCGP487 upon digestion with the restriction endonucleases *Xho*I and *Pst*I. The fragment generated was purified and ligated with  
5 *Xho*I/*Pst*I ends of pBluescript II KS+ (Stratagene, USA). Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP717.

***pCGP716***

10 A 0.95 kb petunia *FLS* terminator fragment downstream from the putative translational stop site was released from the plasmid pCGP487 upon digestion with the restriction endonucleases *Hind*III and *Sac*I. The fragment generated was purified and ligated with *Hind*III/*Sac*I ends of pBluescript II KS+ (Stratagene, USA). Correct insertion of the  
15 fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP716.

**Construction of pCGP493 (short *petFLS* 5':*petFLS*3' expression cassette)**

A 1.8 kb fragment containing the short petunia *FLS* promoter fragment was amplified by PCR using the plasmid pCGP717 as template and the T3 primer (Stratagene, USA) and an  
20 *FLS-Nco* primer (5' AAA ATC GAT ACC ATG GTC TTT TTT TCT TTG TCT ATA C 3') (SEQ ID NO:19). The PCR product was digested with the restriction endonucleases *Xho*I and *Cla*I and the purified fragment was ligated with *Xho*I/*Cla*I ends of pCGP716. Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was  
25 designated as pCGP493.

**Construction of pCGP497 (short *petFLS* 5': *petHfl*: *petFLS*3' expression cassette)**

The petunia *F3'5'H* (*petHfl*) cDNA clone was released from the plasmid pCGP627 (described above) upon digestion with the restriction endonucleases *Bsp*HI and *Fsp*I. The  
30 *Bsp*HI recognition sequence encompasses the putative translation initiating codon and the *Fsp*I recognition sequence commences 2 bp downstream from the stop codon. The petunia

- 68 -

*F3'5'H petHf1* fragment generated was purified and ligated with *Clal* (repaired ends)/*NcoI* ends of the plasmid pCGP493. Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP497.

5

**Construction of pCGP1461 (short *petFLS* 5': *petHf1*: *petFLS*3' binary vector)**

The plasmid pCGP497 was linearised upon digestion with the restriction endonuclease *SacI*. The overhanging ends were repaired and a 4.35 kb fragment containing the *short petFLS* 5': *petHf1*: *petFLS*3' gene expression cassette was released upon digestion with the  
10 restriction endonuclease *KpnI*. The fragment generated was purified and ligated with *PstI* (ends repaired)/*KpnI* ends of the binary vector pWTT2132 (DNAP) (Figure 6). Correct insertion of the fragment in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of pWTT2132 was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting  
15 plasmid was designated as pCGP1461 (Figure 10).

**Plant transformation with pCGP1461**

The T-DNA contained in the binary vector plasmid pCGP1461 (Figure 10) was introduced into rose via *Agrobacterium*-mediated transformation.

20

**The binary vector pCGP1616 (*petRT* 5': *petHf1*: *nos* 3')**

The plasmid pCGP1616 (Figure 11) contains a chimeric petunia *F3'5'H* (*petHf1*) gene under the control of a promoter fragment from the *P. hybrida* 3RT gene (*petRT* 5') (Brugliera, 1994, *supra*) with a terminator fragment from the nopaline synthase gene (*nos* 3') of *Agrobacterium* (Depicker, *et al.*, 1982, *supra*). The chimeric petunia *F3'5'H* cassette  
25 is in a tandem orientation with respect to the 35S 5': *SuRB* gene of the binary vector, pWTT2132 (DNAP) (Figure 6).

- 69 -

*Intermediates in the preparation of the binary vector pCGP1616*

*Isolation of petunia 3RT gene*

*P. hybrida cv. Th7 genomic DNA library construction in EMBL3*

A *Petunia hybrida* cv. Th7 genomic library was prepared according to Sambrook *et al.*  
5 1989, *supra* using a *Sau3A* partial digestion of the genomic DNA. The partially digested  
DNA was cloned into EMBL-3 lambda vector (Stratagene, USA). Screening of the Th7  
genomic library for the petunia 3RT gene was as described in Brugliera, 1994, *supra*.

A 3 kb fragment containing the *petRT* 5': *petHfl*: *nos* 3' cassette was released from the  
10 plasmid pCGP846 (described in Brugliera, 1994, *supra*) upon digestion with the restriction  
endonucleases *Pst*I and *Bam*HI. The purified fragment was ligated with *Pst*I/*Bam*HI ends  
of pWTT2132 (DNAP) (Figure 6). Correct insertion of the fragment in a tandem  
orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette was  
established by restriction endonuclease analysis of plasmid DNA isolated from  
15 tetracycline-resistant transformants. The plasmid was designated as pCGP1616 (Figure  
11).

*Plant transformation with pCGP1616*

The T-DNA contained in the binary vector plasmid pCGP1616 (Figure 11) was introduced  
20 into rose via *Agrobacterium*-mediated transformation.

*The binary vector pCGP1623 (mas/35S: petHfl: ocs 3')*

The plasmid pCGP1623 (Figure 12) contains a chimeric petunia *F3'5'H* (*petHfl*) gene  
under the control of the expression cassette contained in pKIWI101 (Janssen and Gardner,  
25 1989, *supra*) consisting of a promoter fragment from the cauliflower mosaic virus 35S  
gene (35S 5') with an enhancing sequence from the promoter of the mannopine synthase  
gene (*mas*) of *Agrobacterium* and a terminator fragment from the octopine synthase gene  
of *Agrobacterium* (*ocs* 3'). The chimeric petunia *F3'5'H* cassette is in a tandem orientation  
with respect to the 35S 5': *SuRB* gene of the binary vector, pWTT2132 (DNAP) (Figure 6).

30

- 70 -

***Intermediates in the preparation of the binary vector pCGP1623***

The ~1.6 kb fragment of the petunia *F3'5'H petHf1* cDNA clone contained in the plasmid pCGP1303 (Figure 4) was released upon digestion with the restriction endonucleases *BspHI* and *SmaI*. The petunia *F3'5'H petHf1* fragment was purified and ligated with a ~5.9 kb *NcoI/EcoRI* (repaired ends) fragment of pKIWI101 (Janssen and Gardner, 1989, *supra*) to produce the plasmid pCGP1619.

A partial digest of the plasmid pCGP1619 with the restriction endonuclease *XhoI* released a 4.9 kb fragment containing the *mas/35S: petHf1: ocs 3'* expression cassette. The fragment was purified and ligated with *SaII* ends of pWTT2132 (DNAP) (Figure 6). Correct insertion of the fragment in a tandem orientation with respect to the *35S 5': SuRB* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The plasmid was designated as pCGP1623 (Figure 12).

***Plant transformation with pCGP1623***

The T-DNA contained in the binary vector plasmid pCGP1623 (Figure 12) was introduced into rose via *Agrobacterium*-mediated transformation.

***The binary vector pCGP1638 (35S 5': petHf1: ocs 3')***

The plasmid pCGP1638 (Figure 13) contains a chimeric petunia *F3'5'H (petHf1)* gene under the control of a *CaMV 35S* promoter (*35S 5'*) with an octopine synthase terminator (*ocs 3'*). A ~60 bp 5' untranslated leader sequence from the petunia chlorophyll a/b binding protein gene (*Cab 22* gene) (Harpster *et al.*, 1988, *supra*) is included between the *CaMV 35S* promoter fragment and the petunia *F3'5'H petHf1* cDNA clone. The chimeric petunia *F3'5'H* cassette is in a tandem orientation with respect to the *35S 5': SuRB* selectable marker gene cassette of the binary vector, pWTT2132 (Figure 6).

- 71 -

***Intermediates in the preparation of the binary vector pCGP1638******Construction of pCGP1273***

The plasmid pCGP1273 was constructed by subcloning a ~3kb *HindIII/HpaI* fragment containing 35S 5': *GUS: ocs* 3' gene from the binary vector pJJ3499 (Jones *et al.*,  
5 *Transgenic Research*, 1: 285-297, 1992) with the *HindIII/SmaI* ends of the plasmid pBluescript KS II (+) (Stratagene, USA).

***Construction of pCGP1634***

A ~3kb *HindIII/BamHI* fragment containing the 35S 5': *GUS: ocs* 3' gene from  
10 pCGP1273 was then isolated and ligated with the *HindIII/BamHI* ends of the cloning vector pUC19 (New England Biolabs) to create the plasmid pCGP1634.

***Construction of pCGP1636***

The *GUS* fragment from the plasmid pCGP1634 was removed by digesting pCGP1634  
15 with the restriction endonucleases *NcoI* and *XbaI* and purifying the ~3.7kb fragment containing the 35S 5' promoter fragment, the *ocs* 3' terminator fragment and the pUC19 vector backbone.

The petunia *F3'5'H petHf1* cDNA clone was released from pCGP1303 (Figure 4) upon digestion with the restriction endonucleases *BspHI* and *XbaI*. The resulting ~1.6kb  
20 fragment was purified and ligated with the ~3.7kb *NcoI/XbaI* fragment from pCGP1634. Correct insertion of the petunia *F3'5'H petHf1* fragment was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The resulting plasmid containing a 35S 5': *petHf1: ocs* 3' gene was designated pCGP1636.

***Construction of pCGP1638***

The 35S 5': *petHf1: ocs* 3' gene from the plasmid pCGP1636 was released upon digestion of pCGP1636 with the restriction endonucleases *PstI* and *EcoRI*. The ends were repaired and the ~2.6kb fragment was purified and ligated with the *Sam* ends of the binary vector, pWTT2132 (DNAP). Correct insertion of the 35S 5': *petHf1: ocs* 3' gene in a tandem  
30 orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from

- 72 -

tetracycline-resistant transformants. The plasmid was designated as pCGP1638 (Figure 13).

**Plant transformation with pCGP1638**

- 5 The T-DNA contained in the binary vector plasmid pCGP1638 (Figure 13) was introduced into rose via *Agrobacterium*-mediated transformation.

**The binary vector pCGP1860 (RoseCHS 5': petHfl: nos 3')**

- 10 The plasmid pCGP1860 (Figure 14) contains a chimeric petunia *F3'5'H* (*petHfl*) gene under the control of a promoter fragment from the chalcone synthase gene of *Rosa hybrida* (*RoseCHS 5'*) with a terminator fragment from the nopaline synthase gene of *Agrobacterium* (*nos 3'*). The chimeric petunia *F3'5'H* cassette is in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector, pWTT2132 (DNAP) (Figure 6).

15

**Intermediates in the preparation of the binary vector pCGP1860**

**Isolation of Rose CHS promoter**

A rose genomic DNA library was prepared from genomic DNA isolated from young leaves of *Rosa hybrida* cv. Kardinal.

20

The Kardinal genomic DNA library was screened with <sup>32</sup>P-labelled fragment of rose *CHS* cDNA clone contained in the plasmid pCGP634. The rose *CHS* cDNA clone was isolated by screening of a petal cDNA library prepared from RNA isolated from petals of *Rosa hybrida* cv Kardinal (Tanaka *et al.*, 1995, *supra*) using a petunia *CHS* cDNA fragment as probe (clone *1F11* contained in pCGP701, described in Brugliera *et al.*, 1994, *supra*).  
25 Conditions are as described in Tanaka *et al.*, 1995 (*supra*).

30

A rose genomic clone (*roseCHS20N*) was chosen for further analysis and found to contain ~6.4 kb of sequence upstream of the putative initiating methionine of the rose *CHS* coding region.



- 73 -

An ~6.4 kb fragment upstream from the translational initiation site was cloned into pBluescript KS (-) (Statagene) and the plasmid was designated as pCGP1114.

5 The plasmid pCGP1114 was digested with the restriction endonucleases *HindIII* and *EcoRV* to release a 2.7-3.0kb fragment which was purified and ligated with the *HindIII/SmaI* ends of pUC19 (New England Biolabs). Correct insertion of the rose *CHS* promoter fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP1116. The DNA sequence of the rose *CHS* promoter fragment was determined using  
10 pCGP1116 as template (SEQ ID NO:5).

**Construction of pCGP197 (RoseCHS 5': GUS : nos 3' in pUC18 backbone)**

An ~3.0 kb fragment containing the rose chalcone synthase promoter (*RoseCHS 5'*) was released from the plasmid pCGP1116 upon digestion with the restriction endonucleases  
15 *HindIII* and *Asp718*. The fragment was purified and ligated with a *HindIII/Asp718* fragment from pJB1 (Bodeau, 1994, *supra*) containing the vector backbone,  $\beta$ -glucoronidase (*GUS*) and *nos 3'* fragments. Correct insertion of the rose *CHS* promoter fragment upstream of the *GUS* coding sequence was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as  
20 pCGP197.

**Construction of pCGP200 (RoseCHS 5': petHf1: nos 3' in pUC18 backbone)**

A 1.8 kb fragment containing the petunia *F3'5'H* (*petHf1*) fragment was released from the plasmid pCGP1303 (described above) (Figure 4) upon digestion with the restriction  
25 endonucleases *BspHI* and *SacI*. The petunia *F3'5'H petHf1* fragment was purified and ligated with *NcoI/SacI* ends of pCGP197. Correct insertion of the petunia *F3'5'H petHf1* fragment between the rose *CHS* promoter and *nos 3'* fragments was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP200.

30

- 74 -

**Construction of pCGP1860 (RoseCHS 5': petHf1: nos 3' in a binary vector)**

An ~4.9 kb fragment containing the *RoseCHS* 5': *petHf1*: *nos* 3' cassette was released from the plasmid pCGP200 upon digestion with the restriction endonuclease *Bgl*II. The fragment was purified and ligated with *Bam*HI ends of the binary vector, pWTT2132 (DNAP) (Figure 6). Correct insertion of the fragment in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of pWTT2132 was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated as pCGP1860 (Figure 13).

**Plant transformation with pCGP1860**

The T-DNA contained in the binary vector plasmid pCGP1860 (Figure 14) was introduced into rose via *Agrobacterium*-mediated transformation.

**The binary vector pCGP2123 (CaMV 35S: petHf2: ocs 3')**

The plasmid pCGP2123 (Figure 15) contains a chimeric petunia *F3'5'H* (*petHf2*) gene under the control of a CaMV35S promoter with a terminator fragment from the octopine synthase gene of *Agrobacterium* (*ocs* 3'). The chimeric petunia *F3'5'H* cassette is in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector, pCGP1988 (Figure 16).

**Intermediates in the preparation of the binary vector pCGP2123**

**Construction of pCGP1988 (a derivative of the binary vector, pWTT2132)**

The binary vector pCGP1988 (Figure 16) is based on binary vector pWTT2132 (DNAP) (Figure 6) but contains the multi-cloning site from pNEB193 (New England Biolabs). The plasmid pNEB193 was firstly linearized by digestion with the restriction endonuclease *Eco*RI. The overhanging ends were repaired and the multi-cloning fragment was released upon digestion with the restriction endonuclease *Pst*I. The fragment was purified and ligated with *Sa*II (ends repaired)/*Pst*I ends of the binary vector pWTT2132 (DNAP). Correct insertion of the multi-cloning fragment into pWTT2132 was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated as pCGP1988 (Figure 16).

- 75 -

**Construction of pCGP2000 (CaMV 35S promoter fragment in pBluescript)**

The plasmid pCGP2000 was an intermediate plasmid containing a cauliflower mosaic virus (CaMV) 35S promoter fragment in a pBluescript SK (Stratagene, USA) backbone.

- 5 The CaMV 35S promoter fragment from pKIWI101 (Janssen and Gardner, 1989, *supra*) was released upon digestion with the restriction endonucleases *Xba*I and *Pst*I. The ~0.35kb fragment generated was purified and ligated with *Xba*I/*Pst*I ends of the vector pBluescript SK. Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The plasmid was  
10 designated as pCGP2000.

**Construction of pCGP2105 (CaMV 35S 5' and ocs 3' fragments in pBluescript)**

- The plasmid pCGP2105 (Figure 17) contained a CaMV 35S promoter fragment along with a terminator fragment from the octopine synthase gene of *Agrobacterium* (*ocs* 3') both  
15 from pKIWI101 (Janssen and Gardner, 1989, *supra*).

- The *ocs* 3' fragment from pKIWI101 (Janssen and Gardner, 1989, *supra*) was isolated by firstly digesting the plasmid pKIWI101 with the restriction endonuclease *Eco*RI, followed by repair of the overhanging ends, and finally by digestion with the restriction  
20 endonuclease *Xho*I to release a 1.6 kb fragment. This fragment was then ligated with *Hinc*II/*Xho*I ends of pCGP2000. Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The plasmid was designated pCGP2105 (Figure 17).

25 **Construction of pCGP2109 (CaMV 35S: *petHf2*: *ocs* 3' gene in pBluescript)**

The plasmid pCGP2109 contained the CaMV 35S: *petHf2*: *ocs* 3' expression gene cassette in a pBluescript backbone.

- The 1.8 kb petunia *F3'5'H petHf2* cDNA clone was released from pCGP175 (Holton *et al.*,  
30 1993a, *supra*) upon digestion with the restriction endonucleases *Xba*I and *Ssp*I. The overhanging ends were repaired and the purified fragment was ligated with *Pst*I (ends

- 76 -

repaired)/*EcoRV* ends of pCGP2105 (described above) (Figure 17). Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The plasmid was designated pCGP2109.

5     **Construction of pCGP2123 (*CaMV* 35S: *petHf2*: *ocs* 3' binary vector)**

The *CaMV* 35S: *petHf2*: *ocs* 3' cassette was released from pCGP2109 upon digestion with the restriction endonucleases *Asp*718 and *Xba*I. The overhanging ends were repaired and the resultant ~3.7 kb fragment containing the *CaMV* 35S: *petHf2*: *ocs* 3' gene was purified and ligated with repaired ends of *Asp*718 of the binary vector, pCGP1988 (Figure 16).

- 10    Correct insertion of the *CaMV* 35S: *petHf2*: *ocs* 3' gene in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The plasmid was designated as pCGP2123 (Figure 15).

15     **Plant transformation with pCGP2123**

The T-DNA contained in the binary vector plasmid pCGP2123 (Figure 15) was introduced into rose via *Agrobacterium*-mediated transformation.

**EXAMPLE 5**

20     **Analysis of transgenic roses**

The transgenic roses produced in the experiments described in Example 4 were grown to flowering. Flowers were collected and the colors of the petals were coded using the Royal Horticultural Society Colour Charts (RHSCC). The anthocyanins were extracted and the anthocyanidins (specifically the presence of delphinidin or delphinidin-based molecules) analysed by TLC and/or HPLC analysis. Total RNA was also isolated from petal tissue and Northern blot analysis was used to detect transcripts of petunia *F3'5'H* transgenes, endogenous rose *CHS* gene and *SuRB* transgene. The results of the transgenic analysis are summarised in Table 6.

- 77 -

Although over 250 transgenic Kardinal roses were produced (Table 6) none produced flowers with a change in color. TLC and/or HPLC analysis failed to detect accumulation of delphinidin or delphinidin-based molecules pigments confirming the absence of efficient F3'5'H activity. Subsequent Northern analysis on total RNA isolated from petal tissue of these transgenic roses revealed either no detectable intact petunia *F3'5'H* (*petHf1* or *petHf2*) transcripts, or in some cases (see footnotes), degraded transcripts. Hybridization of the same membranes with the selectable marker gene (*SuRB*) or with an endogenous rose *CHS* cDNA probe revealed discrete hybridizing transcripts indicating that the total RNA isolated was not degraded. The detection of the *SuRB* transgene transcripts confirmed that the roses were transgenic.

**TABLE 6** Results of transgenic analysis of rose petals transformed with the T-DNA from various petunia *F3'5'H* (*petHf1* or *petHf2*) gene expression cassettes.

PLASMID	F3'5'H GENE	EVENTS	DEL	RNA
pCGP1452	<i>AmCHS</i> 5': <i>petHf1</i> : <i>petD8</i> 3'	34	0/28	0/34 <sup>1</sup>
pCGP1453	<i>Mac</i> : <i>petHf1</i> : <i>mas</i> 3'	16	0/14	0/13 <sup>2</sup>
pCGP1457	<i>petD8</i> 5': <i>petHf1</i> : <i>petD8</i> 3'	11	0/11	0/11
pCGP1461	<i>short petFLS</i> 5': <i>petHf1</i> : <i>petFLS</i> 3'	11	0/11	0/11
pCGP1616	<i>petRT</i> 5': <i>petHf1</i> : <i>nos</i> 3'	4	0/4	0/4
pCGP1623	<i>mas/35S</i> : <i>petHf1</i> : <i>ocs</i> 3'	27	0/20	0/12 <sup>3</sup>
pCGP1638	<i>CaMV</i> 35S: <i>petHf1</i> : <i>ocs</i> 3'	22	0/14	0/14
pCGP1860	<i>RoseCHS</i> 5': <i>petHf1</i> : <i>nos</i> 3'	15	0/13	0/13
pCGP2123	<i>CaMV</i> 35S: <i>petHf2</i> : <i>ocs</i> 3'	40	0/26	0/10

EVENTS = number of independent transgenic events produced

DEL = number of transgenic events in which delphinidin or delphinidin-based molecules was detected (by TLC or HPLC) in petals over the total number of events analyzed

- 78 -

RNA = number of transgenic events in which intact *F3'5'H* (*petHf1* or *petHf2*) transcripts were detected by Northern blot analysis in total RNA isolated from rose petals over the total number of events analyzed

- 1 = Degraded transcripts were detected in 5 of the 34 analyzed
- 5 2 = Degraded transcripts were detected in 8 of the 13 analyzed
- 3 = Degraded transcripts were detected in 8 of the 12 analyzed

The fact that no intact petunia *F3'5'H* (*petHf1* or *petHf2*) transcripts were ever detected in transgenic rose petals transformed with the T-DNAs described (Table 6) suggested a number of possibilities:

1. that the RNA isolated was degraded. This was not the case as the RNA had been stained by ethidium bromide and visualised under UV-light. The intact visible ribosomal RNA bands were used as an indicator of the quality of the RNA isolated. Furthermore the detection of full-length transcripts of the endogenous rose *CHS* and *SuRB* transgenes confirmed that the RNA preparation was not degraded.

2. that there was no initiation of transcription of the chimeric *F3'5'H* genes evaluated. This was a possibility with some of the expression cassettes analysed, as no *F3'5'H* transcripts were detected by Northern analysis. However all of the petunia *F3'5'H* expression cassettes had proven to be functional (ie. result in an intact transcript and result in the production of delphinidin-based pigments) in other plants such as carnation and petunia.

3. that the petunia *F3'5'H* *petHf1* and *petHf2* mRNAs were unstable in roses. This was also a possibility as degraded petunia *F3'5'H* transcripts were detected by Northern analysis in total RNA isolated from petals of some events. However the petunia *petHf1* and *petHf2* mRNAs had been proven to be stable in other plants such as carnation and petunia. Such instability could be due to aberrant translation leading to mRNA turnover, some feature of the sequence inherently unstable in rose cells, some other factor or factors.

30

- 79 -

There was a need therefore to find suitable promoter fragments that would efficiently drive expression of genes in rose petals and find suitable *F3'5'H* sequences that would result in intact transcripts accumulating in rose petals leading to functional *F3'5'H* activity and to the production of delphinidin-based pigments.

5

**EXAMPLE 6*****Evaluation of promoters in roses*****Development of *GUS* gene expression cassettes.**

The evaluation of the promoter and terminator fragments was performed using the *GUS* reporter gene. Therefore, a number of promoters were linked to the  $\beta$ -glucuronidase reporter gene (*GUS*) (Jefferson *et al.*, 1987, *supra*) and introduced into roses in an attempt to identify expression cassettes that lead to effective initiation of transcription in rose flowers.

15 A summary of the promoters and terminator fragments evaluated is given in Table 7.

**TABLE 7** List of chimaeric *GUS* gene expression cassettes evaluated in roses

PLASMID	<i>GUS</i> EXPRESSION CASSETTE	SELECTABLE MARKER GENE	BACKBONE VECTOR
pCGP1307	<i>petD8 5': GUS: petD8 3'</i>	<i>mas 5': nptII : mas 3'</i>	pCGN1548
pCGP1506	<i>long petFLS 5': GUS: petFLS 3'</i>	<i>nos 5': nptII: nos 3'</i>	pBIN19
pCGP1626	<i>chrysCHS 5': GUS: petRT 3'</i>	<i>35S 5': SuRB</i>	pWTT2132
pCGP1641	<i>petRT 5': GUS: petRT 3'</i>	<i>35S 5': SuRB</i>	pWTT2132
pCGP1861	<i>RoseCHS 5': GUS: nos 3'</i>	<i>35S 5': SuRB</i>	pWTT2132
pCGP1953	<i>AmCHS 5': GUS: petD8 3'</i>	<i>35S 5': SuRB</i>	pWTT2132
pWTT2084	<i>35S 5': GUS: ocs 3'</i>	<i>35S 5': SuRB</i>	pWTT2132

- 80 -

**The binary vector pCGP1307 (*petD8* 5': *GUS*: *petD8* 3')**

The plasmid pCGP1307 (Figure 18) contains a chimeric *GUS* gene under the control of a promoter and terminator fragment from the petunia *PLTP* gene (*petD8* 5' and *petD8* 3', respectively). The chimeric *GUS* reporter gene cassette is in a tandem orientation with respect to the *mas* 5': *nptII*: *mas* 3' selectable marker gene cassette of the binary vector pCGN1548 (McBride and Summerfelt, 1990, *supra*).

**Intermediates in the preparation of the binary vector pCGP1307**

The *nos* 3' fragment from pCGP1101 (see Example 4) was replaced with the 0.75 kb *petD8* 3' fragment (Holton, 1992, *supra*) to produce the plasmid pCGP1106 containing a *petD8* 5': *GUS*: *petD8* 3' expression cassette.

The 5.3 kb fragment containing the *petD8* 5': *GUS*: *petD8* 3' expression cassette was released from the plasmid pCGP1106 upon digestion with the restriction endonucleases *HindIII* and *PstI*. The fragment was purified and ligated with *HindIII*/*PstI* ends of the binary vector, pCGN1548 (McBride and Summerfelt, 1990, *supra*). Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from gentamycin-resistant transformants. The resulting plasmid was designated as pCGP1307 (Figure 18).

**Plant transformation with pCGP1307**

The T-DNA contained in the binary vector plasmid pCGP1307 (Figure 18) was introduced into rose via *Agrobacterium*-mediated transformation.

**The binary vector pCGP1506 (long *petFLS* 5': *GUS*: *petFLS* 3')**

The plasmid pCGP1506 (Figure 19) contains a chimeric *GUS* gene under the control of promoter and terminator fragments from the petunia flavonol synthase gene (*petFLS* 5' and *petFLS* 3', respectively). The chimeric *GUS* reporter gene cassette is in a tandem orientation with respect to the *nos* 5': *nptII*: *nos* 3' selectable marker gene cassette of the binary vector pBIN19 (Bevan, *Nucleic Acids Res* 12: 8711-8721, 1984).



***Intermediates in the preparation of the binary vector pCGP1506***

A 4 kb long petunia *FLS* promoter fragment upstream from the putative translational initiation site was released from the plasmid pCGP486 (described in Example 4) upon digestion with the restriction endonucleases *Xho*I and *Pst*I. The fragment generated was purified and ligated with *Xho*I/*Pst*I ends of pBluescript II KS+ (Stratagene, USA). Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP715.

10 ***Construction of pCGP494 (long petFLS 5':petFLS3' expression cassette)***

A 4.0 kb fragment containing the long petunia *FLS* promoter fragment was amplified by PCR using the plasmid pCGP715 as template and the T3 primer (Stratagene, USA) and an *FLS-Nco* primer (5' AAA ATC GAT ACC ATG GTC TTT TTT TCT TTG TCT ATA C 3') (SEQ ID NO:19). The PCR product was digested with the restriction endonucleases *Xho*I and *Cla*I and the purified fragment was ligated with *Xho*I/*Cla*I ends of pCGP716 (described in Example 4). Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP494.

20 ***Construction of pCGP496 (long petFLS 5': GUS: petFLS3' expression cassette)***

The *GUS* coding sequence from the plasmid pJB1 (Bodeau, 1994, *supra*) was released upon digestion with the restriction endonucleases *Nco*I and *Sma*I. The *GUS* fragment generated was purified and ligated with *Cla*I (repaired ends)/*Nco*I ends of the plasmid pCGP494. Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP496.

30 ***Construction of pCGP1506 (long petFLS 5': GUS: petFLS3' binary vector)***

The plasmid pCGP496 was firstly linearised upon digestion with the restriction endonuclease *Xho*I. The overhanging ends were partially repaired (using only dTTP and dCTP in the reparation reaction) and a 6.7 kb fragment containing the long petFLS 5':

- 82 -

*GUS: petFLS3'* gene expression cassette was released upon digestion with the restriction endonuclease *SacI*. The fragment generated was purified and ligated with *BamHI* (partially repaired ends using dGTP and dATP in the reparation reaction)/*SacI* ends of the binary vector pBIN19. Correct insertion of the fragment in a tandem orientation with respect to the *nos 5': nptII: nos 3'* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from kanamycin-resistant transformants. The resulting plasmid was designated as pCGP1506 (Figure 19).

#### **Plant transformation with pCGP1506**

The T-DNA contained in the binary vector plasmid pCGP1506 (Figure 19) was introduced into rose via *Agrobacterium*-mediated transformation.

#### **The binary vector pCGP1626 (*chrysCHS 5': GUS: petRT 3'*)**

The plasmid pCGP1626 (Figure 20) contains a chimeric *GUS* gene under the control of promoter fragment from the chalcone synthase gene of chrysanthemum (*chrysCHS 5'*) and a terminator fragment from the 3RT gene of petunia (*petRT 3'*) (Brugliera, 1994, *supra*). The chimeric *GUS* reporter gene cassette is in a tandem orientation with respect to the *35S 5': SuRB* selectable marker gene cassette of the binary vector pWTT2132 (DNAP) (Figure 6).

#### **Intermediates in the preparation of the binary vector pCGP1626**

##### **Isolation of chrysanthemum *CHS* promoter**

A chrysanthemum genomic DNA library was prepared from genomic DNA isolated from young leaf material of the chrysanthemum cv Hero.

The chrysanthemum genomic DNA library was screened with <sup>32</sup>P-labelled fragments of a chrysanthemum *CHS* cDNA clone (SEQ ID NO:28) (contained in the plasmid pCGP856) using high stringency conditions. The plasmid pCGP856 contains a 1.5 kb cDNA clone of *CHS* isolated from a petal cDNA library prepared from RNA isolated from the chrysanthemum cv. Dark Pink Pom Pom.

- 83 -

A genomic clone (*CHS5*) was chosen for further analysis and found to contain ~3 kb of sequence upstream of the putative initiating methionine of the chrysanthemum *CHS* coding region.

5 A 4 kb fragment was released upon digestion of the genomic clone *CHS5* with the restriction endonuclease *HindIII*. The fragment containing the chrysanthemum *CHS* promoter was purified and ligated with *HindIII* ends of pBluescript SK (Stratagene, USA). Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was  
10 designated as pCGP1316.

A 2.6 kb chrysanthemum *CHS* promoter fragment upstream from the putative translational initiation site was amplified by PCR using pCGP1316 as template and primers "chrysanCHSATG" (5'-GTTAAGGAAGCCATGGGTGT-3') (SEQ ID NO:8) and the  
15 M13 reverse primer (Stratagene, USA). Primer "chrysanCHSATG" incorporated an *NcoI* restriction endonuclease recognition sequence at the putative translation initiation point for case of cloning. The PCR fragment was purified and ligated with *EcoRV* (dT-tailed) ends of pBluescript KS (Holton and Graham, *Nuc. Acids Res.* 19: 1156, 1990). Correct insertion  
20 of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP1620. The nucleotide sequence of the chrysanthemum *CHS* promoter fragment contained in pCGP1620 is represented as SEQ ID NO:30.

Construction of pCGP1622 (chrysCHS 5': GUS: nos 3' in pUC backbone)

25 A ~2.5 kb fragment containing the chrysanthemum *CHS* promoter was released from the plasmid pCGP1620 upon digestion with the restriction endonucleases *NcoI* and *PstI*. The fragment was purified and ligated with a 4.8 kb *NcoI/PstI* fragment of pJB1 (Bodeau, 1994, *supra*) containing the backbone vector with the *GUS* and *nos 3'* fragments. Correct  
30 insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP1622.

- 84 -

**Construction of pCGP1626 (*chrysCHS* 5': *GUS*: *nos* 3' in binary vector)**

A ~4.6 kb fragment containing the *chrysCHS* 5': *GUS*: *nos* 3' cassette was released from the plasmid pCGP1622 upon digestion with the restriction endonucleases *Pst*I and *Bgl*II.

- 5 The fragment was purified and ligated with *Pst*II/*Bam*HI ends of the binary vector pWTT2132 (DNAP) (Figure 6). Correct insertion of the cassette in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette was established by restriction endonuclease analysis of DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated as pCGP1626 (Figure 20).

10

**Plant transformation with pCGP1626**

The T-DNA contained in the binary vector plasmid pCGP1626 (Figure 20) was introduced into rose via *Agrobacterium*-mediated transformation.

- 15 **The binary vector pCGP1641 (*petRT* 5': *GUS*: *petRT* 3')**

The plasmid pCGP1641 (Figure 21) contains a chimeric *GUS* gene under the control of a petunia 3RT promoter (*petRT* 5') covering 1.1kb upstream from the putative 3RT translation initiation codon with a petunia 3RT terminator (*petRT* 3') covering 2.5 kb downstream from the 3RT stop codon. The chimeric *GUS* cassette is in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector, pWTT2132 (DNAP) (Figure 6).

20

**Intermediates in the preparation of the binary vector pCGP1641**

**Isolation of petunia 3RT gene**

- 25 The isolation of the petunia 3RT gene corresponding to the *Rt* locus of *P. hybrida* has been described in Brugliera, 1994, *supra*.

- 85 -

**Construction of pCGP1625 (CaMV 35S: GUS: petRT 3' cassette)**

The intermediate plasmid pCGP1625 contains a *CaMV 35S: GUS: petRT 3'* cassette in a pUC backbone. The 2.5 kb fragment containing a petRT terminator sequences was released from the plasmid pCGP1610 (described in Brugliera, 1994, *supra*) upon digestion with the restriction endonucleases *Bam*HI and *Sac*I. The fragment was purified and ligated with the *Bgl*III/*Sac*I 4.9kb fragment of pJB1 (Bodeau, 1994, *supra*) containing the vector backbone and the *CaMV 35S* promoter and *GUS* fragments. Correct insertion of the petunia 3RT terminator fragment downstream of the *GUS* fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants.

10 The resulting plasmid was designated as pCGP1625.

**Construction of pCGP1628 (petRT 5': GUS: petRT 3' cassette)**

A 1.1 kb *petRT* promoter fragment was released from the plasmid pCGP1611 (described in Brugliera, 1994, *supra*) upon digestion with the restriction endonucleases *Nco*I and *Pst*I.

15 The purified fragment was ligated with *Nco*I/*Pst*I ends of the 7kb fragment of pCGP1625 containing the vector backbone and the *GUS* and *petRT 3'* fragments. Correct insertion of the *petRT* promoter fragment upstream of the *GUS* fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP1628.

20

**Construction of pCGP1641 (petRT 5': GUS: petRT 3' binary vector)**

A 5.4 kb fragment containing the *petRT 5': GUS: petRT 3'* cassette was released from pCGP1628 upon digestion with the restriction endonuclease *Pst*I. The fragment was purified and ligated with *Pst*I ends of the binary vector pWTT2132 (DNAP) (Figure 6).

25 Correct insertion of the fragment in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated as pCGP1641 (Figure 21).

- 86 -

**Plant transformation with pCGP1641**

The T-DNA contained in the binary vector plasmid pCGP1641 (Figure 21) was introduced into rose via *Agrobacterium*-mediated transformation.

**5 The binary vector pCGP1861 (RoseCHS 5': GUS: nos 3')**

The plasmid pCGP1861 (Figure 22) contains a chimeric *GUS* gene under the control of a promoter fragment from the *CHS* gene of *R. hybrida* (*RoseCHS 5'*) with a terminator fragment from the *nos* gene of *Agrobacterium* (*nos 3'*). The chimeric *GUS* reporter gene cassette is in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector, pWTT2132 (Figure 6).

An ~5 kb fragment containing the *RoseCHS 5': GUS: nos 3'* cassette was released from pCGP197 (described in Example 4) upon digestion with the restriction endonuclease *Bgl*II. The fragment was purified and ligated with *Bam*HI ends of the binary vector, pWTT2132 (DNAP). Correct insertion of the fragment in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated as pCGP1861 (Figure 22).

**20 Plant transformation with pCGP1861**

The T-DNA contained in the binary vector plasmid pCGP1861 (Figure 22) was introduced into rose via *Agrobacterium*-mediated transformation.

**The binary vector pCGP1953 (AmCHS 5': GUS: petD8 3')**

25 The plasmid pCGP1953 (Figure 23) contains a chimeric *GUS* gene under the control of a promoter fragment from the *CHS* gene of *Antirrhinum majus* (*AmCHS 5'*) with a petunia *PLTP* terminator (*petD8 3'*). The chimeric *GUS* reporter gene cassette is in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector, pWTT2132 (DNAP) (Figure 6).

30

- 87 -

***Intermediates in the preparation of the binary vector pCGP1953***

The plasmid pJB1 (Bodeau, 1994, *supra*) was linearised with the restriction endonuclease *Nco*I. The overhanging ends were repaired and the 1.8 kb *GUS* fragment was released upon digestion with *Bam*HI. The *GUS* fragment was purified and was ligated with the 5 kb *Xba*I(ends repaired)/*Bam*HI fragment of pCGP726 containing the pBluescript backbone vector and the *AmCHS* 5' and *petD8* 3' fragments (described in Example 4). Correct insertion of the *GUS* fragment between the *AmCHS* 5' and *petD8* 3' fragments was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The plasmid was designated as pCGP1952.

A 3.8 kb fragment containing the *AmCHS* 5': *GUS*: *petD8* 3' expression cassette was released from the plasmid pCGP1952 upon digestion with the restriction endonucleases *Eag*I and *Pst*I. The overhanging ends were repaired and the purified fragment was ligated with the repaired ends of an *Asp*718 digested pWTT2132 binary vector (Figure 6). Correct insertion of the fragment in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The plasmid was designated as pCGP1953 (Figure 23).

***Plant transformation with pCGP1953***

The T-DNA contained in the binary vector plasmid pCGP1953 (Figure 23) was introduced into rose via *Agrobacterium*-mediated transformation.

***The binary vector pWTT2084 (35S 5': GUS: ocs 3')***

The plasmid pWTT2084 (DNAP) (Figure 24) contains a chimeric *GUS* gene under the control of a *CaMV* 35S promoter (35S 5') with an octopine synthase terminator (*ocs* 3'). An ~60 bp 5' untranslated leader sequence from the petunia chlorophyll a/b binding protein gene (*Cab* 22 gene) (Harpster *et al.*, 1988, *supra*) is included between the *CaMV* 35S promoter fragment and the *GUS* clone. The chimeric *GUS* cassette is in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector, pWTT2084.

**Plant transformation with pWTT2084**

The T-DNA contained in the binary vector plasmid pWTT2084 (Figure 24) was introduced into rose via *Agrobacterium*-mediated transformation.

5

Transgenic analysis of roses transformed with GUS expression cassettes

Northern blot analysis was performed on total RNA isolated from petals of developmental stages 3 to 4 of transgenic Kardinal roses transformed with the T-DNA of various *GUS* expression cassettes. There was either no accumulating transcript or an intact full-length transcript of the expected size of ~1.8kb as detected by Northern blot hybridisation. The relative levels of *GUS* transcripts accumulating in the rose petals were recorded (see Table 8).

10

**TABLE 8** Summary of Northern analysis on transgenic Kardinal rose flowers (open bud stage) containing *GUS* constructs.

15

PLASMID	<i>GUS</i> REPORTERGENE	SELECTABLE MARKER GENE	<i>GUS</i> TRANSCRIPT LEVELS
pCGP1307	<i>petD8</i> 5': <i>GUS</i> : <i>petD8</i> 3'	<i>mas</i> 5': <i>nptII</i> : <i>mas</i> 3'	—
pCGP1506	<i>petFLS</i> 5': <i>GUS</i> : <i>petFLS</i> 3'	<i>nos</i> 5': <i>nptII</i> : <i>nos</i> 3'	—
pCGP1626	<i>chrysCHS</i> 5': <i>GUS</i> : <i>petRT</i> 3'	<i>35S</i> 5': <i>SuRB</i>	++ to +++
pCGP1641	<i>petRT</i> 5': <i>GUS</i> : <i>petRT</i> 3'	<i>35S</i> 5': <i>SuRB</i>	—
pCGP1861	<i>RoseCHS</i> 5': <i>GUS</i> : <i>nos</i> 3'	<i>35S</i> 5': <i>SuRB</i>	++++
pCGP1953	<i>AmCHS</i> 5': <i>GUS</i> : <i>petD8</i> 3'	<i>35S</i> 5': <i>SuRB</i>	—
pWTT2084	<i>35S</i> 5': <i>GUS</i> : <i>ocs</i> 3'	<i>35S</i> 5': <i>SuRB</i>	+++++

— = no transcripts detected

+ to +++++ = relative levels (low to high) of full-length *GUS* transcript detected by Northern blot analysis

20



- 89 -

Based on the above results (Table 8), the *CaMV 35S* (*35S 5'*) and rose *CHS* (*RoseCHS 5'*) promoters appear to drive relatively high levels of transcription in rose petals. The chrysanthemum *CHS* promoter (*chrysCHS 5'*) appears to also lead to high transcript levels but not as high as those obtained using *CaMV 35S* or rose *CHS* promoters. Surprisingly, antirrhinum (snapdragon) *CHS* (*AmCHS 5'*), petunia *3RT* (*petRT 5'*), petunia *FLS* (*petFLS 5'*) and petunia *PLTP* - (*petD8 5'*) promoters did not appear to function in rose petals as no *GUS* transcripts were detected with expression cassettes incorporating these promoters. However, these same promoters fused to *petHf1* and/or -*GUS* genes had previously been proven to function well in carnation and petunia leading to relatively high full-length transcript levels and for *petHf1* genes, the production of delphinidin or delphinidin-based molecules pigments. The result obtained with the antirrhinum *CHS* promoter (*AmCHS 5'*) fused with the *GUS* gene was more surprising as promoter regions from homologous genes from two other species (rose and chrysanthemum) appeared to function relatively well in roses. The antirrhinum *CHS* promoter had also been successfully used in conjunction with petunia *F3'5'H* (*petHf1*) to produce the novel violet-colored carnations Florigene Moondust (see International Patent Application No. PCT/AU96/00296).

The evaluation of promoter and terminator fragments fused with the *GUS* gene also provided further evidence to suggest that the petunia *F3'5'H* *petHf1* and *petHf2* sequences were unstable in roses as constructs containing the petunia *F3'5'H* sequences ligated to the *CaMV 35S*, -rose *CHS* and chrysanthemum *CHS* promoters (which do function in rose) did not result in intact petunia *F3'5'H* *petHf1* or *petHf2* transcripts in roses (see Table 6).

#### EXAMPLE 7

##### 25 *Isolation of F3'5'H sequences from species other than petunia*

Since the petunia *F3'5'H* sequences had already been proven to function in various plants such as carnation, petunia and tobacco and ultimately resulted in the production of delphinidin-based pigments, it was reasonable to assume that these sequences would also prove functional in roses. There was an assumption that the enzyme activity may vary depending on the background of the species, indeed between cultivars of a given species, that the petunia *F3'5'H* was introduced into. However, there was no expectation that full-

- 90 -

length recombinant petunia *F3'5'H* mRNA would not accumulate. Analysis of the petunia *F3'5'H* nucleotide sequences (*petHf1* and *petHf2*) did not reveal any sequences which might lead to instability and subsequent degradation (Johnson *et al.*, *In A look beyond transcription*, ASPP, USA, Bailey-Serres and Gallie, eds, 1998), intron: exon splice  
5 junctions (Brendel *et al.*, *In A look beyond transcription*, ASPP, USA, Bailey-Serres and Gallie, eds, 1998), or any autocatalytic or degradation trigger sequences reported in the scientific literature to date (*In A look beyond transcription*, ASPP, USA, Bailey-Serres and Gallie, eds, 1998). The surprising result suggested that there were factors specific to rose that resulted in petunia *F3'5'H* sequences being unstable.

10

Since it was not obvious why the petunia *F3'5'H* sequences were unstable in roses but stable in carnation, petunia or tobacco, a number of *F3'5'H* sequences were isolated across a range of families in an attempt to determine whether any *F3'5'H* sequence would be  
15 stable *F3'5'H* transcripts and *F3'5'H* activity and ultimately the production of delphinidin-based pigments in roses leading to a change in flower color.

#### Construction of petal cDNA libraries

Petal cDNA libraries were prepared from RNA isolated from petals from bud to opened  
20 flower stages from various species of plants described in Table 9. *Rosa hybrida* is classified in the family Rosaceae, Order Rosales, Subclass Rosidae and so species that produced delphinidin-based pigments and so contained a functional *F3'5'H* and belonged to the Subclass Rosidae were selected. *Petunia hybrida* is classified in the Family Solanaceae, Order Solanales, Subclass Asteridae and so species from the Subclass  
25 Asteridae that produced delphinidin-based pigments were also selected.

**TABLE 9** List of flowers from which total RNA was isolated for the preparation of petal cDNA libraries. Information obtained from National Center for Biotechnology Information (NCBI) website under Taxonomy browser (TaxBrowser) as of August 2003.

FLOWER	SPECIES	FAMILY	ORDER	SUBCLASS
gentian	<i>Gentiana spp.</i>	Gentianaceae	Gentianales	Asteridae
lavender	<i>Lavandula spp.</i>	Lamiaceae	Lamiales	Asteridae
salvia	<i>Salvia spp.</i>	Lamiaceae	Lamiales	Asteridae
sollya	<i>Sollya spp.</i>	Pittosporaceae	Apiales	Asteridae
kennedia	<i>Kennedia spp.</i>	Fabaceae	Fabales	Rosidae
butterfly pea	<i>Clitoria ternatea</i>	Fabaceae	Fabales	Rosidae
pansy	<i>Viola spp.</i>	Violaceae	Malpighiales	Rosidae

Unless otherwise described, total RNA was isolated from the petal tissue of purple/blue flowers using the method of Turpen and Griffith (*BioTechniques* 4: 11-15, 1986). Poly(A)<sup>+</sup> RNA was selected from the total RNA, using oligotex-dT<sup>TM</sup> (Qiagen) or by three cycles of oligo-dT cellulose chromatography (Aviv and Leder, *Proc. Natl. Acad. Sci. USA* 69: 1408, 1972).

In general a  $\lambda$ ZAPII/ Gigapack II Cloning kit (Stratagene, USA) (Short *et al.*, *Nucl. Acids Res.* 16: 7583-7600, 1988) was used to construct directional petal cDNA libraries in  $\lambda$ ZAPII using around 5  $\mu$ g of poly(A)<sup>+</sup> RNA isolated from petal as template. The total number of recombinants obtained was generally in the order of  $1 \times 10^5$  to  $1 \times 10^6$ .

After transfecting XL1-Blue MRF' cells, the packaged cDNA mixtures were plated at around 50,000 pfu per 15 cm diameter plate. The plates were incubated at 37°C for 8 hours, and the phage were eluted in 100 mM NaCl, 8 mM MgSO<sub>4</sub>, 50 mM Tris-HCl pH 8.0, 0.01% (w/v) gelatin (Phage Storage Buffer (PSB)) (Sambrook *et al.*, 1989, *supra*). Chloroform was added and the phages stored at 4°C as amplified libraries.

- 92 -

In general around 100,000 pfu of the amplified libraries were plated onto NZY plates (Sambrook *et al.*, 1989, *supra*) at a density of around 10,000 pfu per 15 cm plate after transfecting XL1-Blue MRF' cells, and incubated at 37°C for 8 hours. After incubation at 4°C overnight, duplicate lifts were taken onto Colony/Plaque Screen™ filters (DuPont) and  
5 treated as recommended by the manufacturer.

#### Plasmid Isolation

Helper phage R408 (Stratagene, USA) was used to excise pBluescript phagemids containing cDNA inserts from amplified λZAPII or λZAP cDNA libraries using methods  
10 described by the manufacturer.

#### Screening of petal cDNA Libraries

Prior to hybridization, duplicate plaque lifts were washed in prewashing solution (50 mM Tris-HCl pH7.5, 1 M NaCl, 1 mM EDTA, 0.1% (w/v) sarcosine) at 65°C for 30 minutes;  
15 followed by washing in 0.4 M sodium hydroxide at 65°C for 30 minutes; then washed in a solution of 0.2 M Tris-HCl pH 8.0, 0.1 x SSC, 0.1% (w/v) SDS at 65°C for 30 minutes and finally rinsed in 2 x SSC, 1.0% (w/v) SDS.

The membrane lifts from the petal cDNA libraries were hybridized with <sup>32</sup>P-labelled  
20 fragments of a 1.6 kb *Bsp*HI/*Fsp*I fragment from pCGP602 (Figure 2) (SEQ ID NO: 1) containing the petunia *F3'5'H petHf1* cDNA clone (Holton *et al.*, 1993a, *supra*).

Hybridization conditions included a prehybridization step in 10% v/v formamide, 1 M NaCl, 10% w/v dextran sulphate, 1% w/v SDS at 42°C for at least 1 hour. The <sup>32</sup>P-labelled  
25 fragments (each at 1x10<sup>6</sup> cpm/mL) were then added to the hybridization solution and hybridization was continued at 42°C for a further 16 hours. The filters were then washed in 2 x SSC, 1% w/v SDS at 42°C for 2 x 1 hour and exposed to Kodak XAR film with an intensifying screen at -70°C for 16 hours.

- 93 -

Strongly hybridizing plaques were picked into PSB (Sambrook *et al.*, 1989, *supra*) and rescreened to isolate purified plaques, using the plating and hybridization conditions as described for the initial screening of the cDNA library. The plasmids contained in the  $\lambda$ ZAPII or  $\lambda$ ZAP bacteriophage vectors were rescued and sequence data was generated from the 3' and 5' ends of the cDNA inserts. New *F3'5'H* cDNA clones were identified based on sequence similarity to the petunia *F3'5'H petHf1* cDNA clone.

The cDNA clones isolated were given plasmid designation numbers as described in Table 10.

**TABLE 10** Plasmid numbers and SEQ ID NO. of *F3'5'H* cDNA clones isolated from various species

SPECIES	CLONE	PLASMID NUMBER	FIGURE NUMBER	SEQ ID NO.
<i>Viola spp.</i>	BP#18	pCGP1959	25	9
<i>Viola spp.</i>	BP#40	pCGP1961	26	11
<i>Salvia spp.</i>	Sal#2	pCGP1995	31	13
<i>Salvia spp.</i>	Sal#47	pCGP1999	32	15
<i>Sollya spp.</i>	Soll#5	pCGP2110	37	17
<i>Kennedia spp.</i>	Kenn#31	pCGP2231	40	26
<i>Clitoria ternatea</i>	BpeaHF2	pBHF2F4	43	20
<i>Gentiana triflora</i>	Gen#48	pG48	47	22
<i>Lavandula nil</i>	LBG	pLHF8	51	31

#### 15 Viola (pansy) *F3'5'H* constructs

##### Isolation of *F3'5'H* cDNA clones from petals of *Viola spp.* (pansy)

Total RNA and poly (A)<sup>+</sup> RNA was isolated from petals of young buds of *Viola spp.* cultivar black pansy as described above. A petal cDNA library was constructed using  $\lambda$ ZAPII/ Gigapack II Cloning kit (Stratagene, USA) and screened as described above. Two full-length pansy *F3'5'H* cDNA clones (*BP#18* (SEQ ID NO:9) in pCGP1959 (Figure 25)

- 94 -

and BP#40 (SEQ ID NO:11) in pCGP1961 (Figure 26)) were identified by sequence similarity to the petunia *F3'5'H petHf1* cDNA clone (SEQ ID NO:1). The BP#18 and BP#40 shared 82% identity at the nucleotide level. Comparison of the nucleotide sequence of pansy *F3'5'H* clones (BP#18 and BP#40) with that of the petunia *F3'5'H* revealed  
5 around 60% identity to the petunia *F3'5'H petHf1* clone and 62% identity to the petunia *F3'5'H petHf2* clone.

The binary vectors, pCGP1972 and pCGP1973 (*AmCHS* 5': BP#18 or BP#40: *petD8* 3')  
The plasmids pCGP1972 (Figure 27) and pCGP1973 (Figure 28) contain the pansy *F3'5'H*  
10 cDNA clone (BP#18 and BP#40, respectively) between an *A. majus* (snapdragon) *CHS* promoter fragment (*AmCHS* 5') and a petunia *PLTP* terminator fragment (*petD8* 3'). The chimeric *F3'5'H* genes are in tandem with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector, pWTT2132 (DNAP) (Figure 6).

15 The petunia *F3'5'H (petHf1)* cDNA clone in pCGP725 (described in Example 4) (Figure 7) was removed by initially digesting pCGP725 with the restriction endonuclease *Bam*HI. The ends were repaired and the linearised plasmid was further digested with the restriction endonuclease *Xba*I. The ~4.9kb fragment containing the vector with the *AmCHS* 5' and *petD8* 3' fragments was purified and ligated with the ~1.6kb *Kpn*I (ends repaired)/*Xba*I  
20 fragment containing the pansy *F3'5'H* BP#18 or BP#40 cDNA clone from pCGP1959 or pCGP1961, respectively to produce pCGP1970 and pCGP1971, respectively. The *AmCHS* 5': pansy *F3'5'H*: *petD8* 3' cassette was then isolated from pCGP1970 or pCGP1971 by firstly digesting with the restriction endonuclease *Not*I. The ends of the linearised plasmid were repaired and then the chimeric *F3'5'H* genes were released upon digestion with the  
25 restriction endonuclease *Eco*RV. The purified fragments were then ligated with *Asp*718 (repaired ends) of the binary vector pWTT2132 (DNAP). Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmids were designated pCGP1972 (Figure 27) and pCGP1973 (Figure 28), respectively.

30

- 95 -

**Carnation and petunia transformation with pCGP1972 and 1973**

The T-DNAs contained in the binary vector plasmids pCGP1972 (Figure 27) and pCGP1973 (Figure 28) were introduced separately into *Dianthus caryophyllus* cultivars Kortina Chanel and Monte Lisa and *Petunia hybrida* cv. Skr4 x Sw63 via *Agrobacterium*-mediated transformation.

**The binary vectors, pCGP1967 and pCGP1969 (CaMV 35S: pansy F3'5'H: ocs 3')**

The binary vectors pCGP1967 (Figure 29) and pCGP1969 (Figure 30) contain chimeric *CaMV* 35S: *pansy* F3'5'H: *ocs* 3' genes in tandem with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector, pWTT2132 (DNAP) (Figure 6).

**Intermediates in the preparation of the binary vectors pCGP1967 and pCGP1969**

The plasmids pCGP1959 (Figure 25) and pCGP1961 (Figure 26) were firstly linearized upon digestion with the restriction endonuclease *Kpn*I. The overhanging *Kpn*I ends were repaired and the *pansy* F3'5'H cDNA clones, BP#18 and BP#40, were released upon digestion with the restriction endonuclease *Pst*I. The ~1.6 kb fragments generated were ligated with an ~5.9 kb *Eco*RI (repaired ends)/*Pst*I fragment of pKIWI101 (Janssen and Gardner, 1989, *supra*). Correct insertion of each fragment was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The resulting plasmids were designated pCGP1965 and pCGP1966, respectively.

The plasmids pCGP1965 and pCGP1966 were firstly partially digested with the restriction endonuclease *Xho*I. The resulting fragments were further digested with the restriction endonuclease *Xba*I. The overhanging ends were repaired and the 3.6kb fragments containing the *CaMV* 35S: *pansy* F3'5'H: *ocs* 3' chimeric genes were isolated and ligated with *Asp*718 repaired ends of pWTT2132 (Figure 6). Correct insertion of each fragment was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmids were designated pCGP1967 (Figure 29) and pCGP1969 (Figure 30), respectively.

- 96 -

**Rose transformation with pCGP1967 and pCGP1969**

The T-DNAs contained in the binary vector plasmids pCGP1967 (Figure 29) and pCGP1969 (Figure 31) were introduced separately into *Rosa hybrida* cv. Kardinal and Soft Promise via *Agrobacterium*-mediated transformation. The T-DNA contained in the binary

5 vector plasmids pCGP1969 (Figure 31) was also introduced into *Rosa hybrida* cv. Pamela and Medeo via *Agrobacterium*-mediated transformation.

**Salvia F3'5'H constructs****Isolation of a F3'5'H cDNA clone from petals of Salvia spp.**

10 Total RNA and poly (A)<sup>+</sup> RNA was isolated from young petal buds of *Salvia* spp. (bought from a nursery) as described above. A petal cDNA library was constructed using λZAPII/ Gigapack II Cloning kit (Stratagene, USA). Two full-length salvia F3'5'H cDNA clones (Sal#2 (SEQ ID NO:13) in pCGP1995 (Figure 31) and Sal#47 (SEQ ID NO:15) in pCGP1999 (Figure 32)) were identified by sequence similarity with the petunia F3'5'H

15 *petHf1* cDNA clone. The Sal#2 and Sal#47 shared 95% identity at the nucleotide level. Comparison of the nucleotide sequence of salvia F3'5'H clones (Sal#2 and Sal#47) with that of the petunia F3'5'H revealed around 57% identity to the petunia F3'5'H *petHf1* clone (SEQ ID NO:1) and 58% identity to the petunia F3'5'H *petHf2* clone (SEQ ID NO:3).

20

**The binary vectors, pCGP2121 and pCGP2122*****(AmCHS 5': Salvia F3'5'H #2 or #47: petD8 3')***

The plasmids pCGP2121 (Figure 33) and pCGP2122 (Figure 34) contain the salvia F3'5'H cDNA clones (Sal#2 and Sal#47, respectively) between a snapdragon *CHS* promoter

25 fragment (*AmCHS* 5') and a petunia *PLTP* terminator fragment (*petD8* 3') in tandem with the 35S 5': *SuRB* selectable marker gene cassette of the binary vector pWTT2132 (DNAP) (Figure 6).



- 97 -

The petunia *F3'5'H* (*petHf1*) cDNA clone in pCGP725 (described in Example 4) (Figure 7) was removed by initially digesting pCGP725 with the restriction endonuclease *Bam*HI. The ends were repaired and the linearised plasmid was further digested with the restriction endonuclease *Xba*I. The ~4.9kb fragment containing the vector with the *AmCHS* 5' and  
 5 *petD8* 3' fragments was purified and ligated with the ~1.6kb *Xho*I/*Bam*HI (ends repaired) fragment from pCGP1995 (Figure 31) containing the salvia *F3'5'H* #2 or *Xho*I/*Eco*RI (ends repaired) fragment from pCGP1999 (Figure 32) containing the salvia *F3'5'H* #47, respectively to produce pCGP2116 and pCGP2117, respectively.

10 The *AmCHS* 5': salvia *F3'5'H*: *petD8* 3' cassette was isolated from pCGP2116 or pCGP2117 by firstly digesting with the restriction endonuclease *Not*I. The ends of the linearized plasmid were repaired and then the chimeric *F3'5'H* gene cassettes were released upon digestion with the restriction endonuclease *Eco*RV. The ~3.6kb purified fragments  
 15 (described in Example 4). Correct insertion of each fragment was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmids were designated pCGP2121 (Figure 33) and pCGP2122 (Figure 34), respectively.

#### 20 *Carnation and petunia transformation with pCGP2121 and pCGP2122*

The T-DNAs contained in the binary vector plasmids pCGP2121 (Figure 33) and pCGP2122 (Figure 34) were introduced separately into *Dianthus caryophyllus* cultivars Kortina Chanel and Monte Lisa and *Petunia hybrida* cv. Skr4 x Sw63 via *Agrobacterium*-mediated transformation.

25

#### *The binary vectors, pCGP2120 and pCGP2119 (CaMV 35S: salvia F3'5'H: ocs 3')*

The binary vectors pCGP2120 (Figure 35) and pCGP2119 (Figure 36) contain chimeric *CaMV* 35S: salvia *F3'5'H*: *ocs* 3' gene cassettes in tandem with the 35S 5': *SuRB* selectable marker gene cassette of the binary vector pCGP1988 (Figure 16).

30

- 98 -

***Intermediates in the preparation of the binary vectors pCGP2120 and pCGP2119***

The plasmids pCGP1995 (Figure 31) and pCGP1999 (Figure 32) were firstly linearized upon digestion with the restriction endonuclease *Xho*I. The overhanging *Xho*I ends were repaired and then the salvia *F3'5'H* cDNA clones Sal#2 or Sal#47 were released upon digestion with the restriction endonuclease *Eco*RI. In the case of pCGP1995 a partial digest with *Eco*RI was undertaken. The ~1.7 kb fragments were ligated with the *Cla*I (repaired ends)/*Eco*RI ends of pCGP2105 (Figure 17). Correct insertion of each fragment was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The resulting plasmids were designated pCGP2112 and pCGP2111, respectively.

The plasmids pCGP2112 and pCGP2111 were digested with the restriction endonucleases *Xho*I and *Xba*I. The resulting overhanging ends were repaired and ~3.6 kb fragments containing the *CaMV 35S: salvia F3'5'H: ocs 3'* chimeric genes were isolated and ligated with *Asp*718 repaired ends of the binary vector, pCGP1988 (described in Example 4). Correct insertion of each fragment was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmids were designated pCGP2120 (Figure 35) and pCGP2119 (Figure 36), respectively.

***Rose transformation with pCGP2120 and pCGP2119***

The T-DNAs contained in the binary vector plasmids pCGP2120 (Figure 35) and pCGP2119 (Figure 36) were introduced separately into *Rosa hybrida* cv. Kardinal via *Agrobacterium*-mediated transformation.

***Sollya F3'5'H constructs******Isolation of a F3'5'H cDNA clone from petals of Sollya spp.***

Total RNA and poly (A)<sup>+</sup> RNA was isolated from young petal buds of *Sollya* spp. (bought from a nursery) as described above. A petal cDNA library was constructed using λZAPII/Gigapack II Cloning kit (Stratagene, USA). One full-length *Sollya F3'5'H* cDNA clone (Sol#5 (SEQ ID NO:17) in pCGP2110 (Figure 37)) was identified by sequence similarity to the petunia *F3'5'H petHf1* cDNA clone. Comparison of the nucleotide sequence of the

- 99 -

sollya *F3'5'H* clone with that of the petunia *F3'5'H* revealed around 48% identity to the petunia *F3'5'H petHf1* clone (SEQ ID NO:1) and 52% identity to the petunia *F3'5'H petHf2* clone (SEQ ID NO:3).

5 **The binary vector pCGP2130 (*AmCHS* 5': Sollya *F3'5'H*: *petD8* 3')**

The plasmid pCGP2130 (Figure 38) contains the sollya *F3'5'H* Soll#5 cDNA clone between a snapdragon *CHS* promoter fragment (*AmCHS* 5') and a petunia *PLTP* terminator fragment (*petD8* 3') in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector pCGP1988 (Figure 16).

10

The petunia *F3'5'H (petHf1)* cDNA clone in pCGP725 (described in Example 4) (Figure 7) was removed by initially digesting pCGP725 with the restriction endonucleases *XbaI* and *BamHI*. The ends were repaired the ~4.9kb fragment containing the vector with the *AmCHS* 5' and *petD8* 3' fragments was purified and ligated with the repaired ends of the  
15 ~1.6kb *Asp718/PstI* fragment from pCGP2110 containing the sollya *F3'5'H* cDNA clone to produce pCGP2128. Correct insertion of the sollya *F3'5'H* fragment in tandem with the *AmCHS* 5' and *petD8* 3' fragments was confirmed by restriction endonuclease mapping.

20

The *AmCHS* 5': sollya *F3'5'H*: *petD8* 3' gene cassette was then isolated from pCGP2128 by firstly digesting with the restriction endonuclease *NotI*. The ends of the linearized plasmid were repaired and then the chimeric *F3'5'H* gene was released upon digestion with the restriction endonuclease *EcoRV*. The ~3.5kb purified fragment was then ligated with *Asp718* (repaired ends) of the binary vector pCGP1988 (described in Example 4) (Figure 16). Correct insertion of the fragment was established by restriction endonuclease analysis  
25 of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated pCGP2130 (Figure 38).

**Carnation and petunia transformation with pCGP2130**

30

The T-DNA contained in the binary vector plasmid pCGP2130 (Figure 38) was introduced into *Dianthus caryophyllus* cultivars Kortina Chamel and Monte Lisa and *Petunia hybrida* cv. Skr4 x Sw63 via *Agrobacterium*-mediated transformation.

- 100 -

**The binary vector pCGP2131 (*CaMV 35S: sollya F3'5'H: ocs 3'*)**

The binary vector pCGP2131 (Figure 39) contains a chimeric *CaMV 35S: sollya F3'5'H: ocs 3'* gene in tandem with the 35S 5': *SuRB* selectable marker gene cassette of the binary  
5 vector pCGP1988 (Figure 16).

**Intermediates in the preparation of the binary vector pCGP2131**

The plasmid pCGP2110 was firstly linearized upon digestion with the restriction endonuclease *Asp718*. The overhanging ends were repaired and then the *sollya F3'5'H*  
10 cDNA clone was released upon digestion with the restriction endonuclease *PstI*. The ~1.7 kb fragment was ligated with the *EcoRV/PstI* ends of pCGP2105 (Figure 17). Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated pCGP2129.

15 A 3.6 kb fragment containing the *CaMV 35S: sollya F3'5'H: ocs 3'* chimeric gene was released upon digestion with the restriction endonucleases *Asp718* and *XbaI*. The overhanging ends were repaired and the purified fragment was ligated with of *Asp718* repaired ends of the binary vector, pCGP1988 (Figure 16). Correct insertion of the  
20 fragment was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated pCGP2131 (Figure 39).

**Rose transformation with pCGP2131**

25 The T-DNA contained in the binary vector plasmid pCGP2131 (Figure 39) was introduced into *Rosa hybrida* cv. Kardinal via *Agrobacterium*-mediated transformation.

- 101 -

**Kennedia F3'5'H constructs****Isolation of a F3'5'H cDNA clone from petals of *Kennedia* spp.**

Total RNA and poly (A)<sup>+</sup> RNA was isolated from young petal buds of *Kennedia* spp. (bought from a nursery) as described above. A petal cDNA library was constructed using

5  $\lambda$ ZAPII/ Gigapack II Cloning kit (Stratagene, USA). One full-length kennedia F3'5'H cDNA clone (*Kenn*#31 in pCGP2231 (Figure 40)) (SEQ ID NO:26) was identified by sequence similarity to the petunia F3'5'H *petHf1* cDNA clone. Comparison of the nucleotide sequence of the kennedia F3'5'H clone with that of the petunia F3'5'H revealed around 64% identity to the petunia F3'5'H *petHf1* clone (SEQ ID NO:1) and 60% identity

10 to the petunia F3'5'H *petHf2* clone (SEQ ID NO:3).

**The binary vector pCGP2256 (*AmCHS* 5': kennedia F3'5'H: *petD8* 3')**

The plasmid pCGP2256 (Figure 41) contains the kennedia F3'5'H (*Kenn*#31) cDNA clone between a snapdragon *CHS* promoter fragment (*AmCHS* 5') and a petunia *PLTP* terminator

15 fragment (*petD8* 3') in tandem with the 35S 5': *SuRB* selectable marker gene cassette of the binary vector pCGP1988 (Figure 16).

The petunia F3'5'H (*petHf1*) cDNA clone in pCGP725 (described in Example 4) (Figure 7) was removed by initially digesting pCGP725 with the restriction endonucleases *Xba*I and

20 *Bam*HI. The ends were repaired the ~4.9kb fragment containing the vector with the *AmCHS* 5' and *petD8* 3' fragments was purified and ligated with the repaired ends of the ~1.8kb *Xho*I/*Bam*HI fragment from pCGP2231 containing the kennedia F3'5'H cDNA clone to produce pCGP2242. Correct insertion of the kennedia F3'5'H fragment in tandem with the *AmCHS* 5' and *petD8* 3' fragments was confirmed by restriction endonuclease

25 mapping.

The *AmCHS* 5': kennedia F3'5'H: *petD8* 3' cassette was then isolated from pCGP2242 by digesting with the restriction endonucleases *Not*I and *Eco*RI. The ends were repaired and the ~3.7kb purified fragment was then ligated with *Asp*718 repaired ends of the binary

30 vector pCGP1988 (described in Example 4) (Figure 16). Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from

- 102 -

tetracycline-resistant transformants. The resulting plasmid was designated pCGP2256 (Figure 41).

**Petunia transformation with pCGP2256**

- 5 The T-DNA contained in the binary vector plasmid pCGP2256 (Figure 41) was introduced into *Petunia hybrida* cv. Skr4 x Sw63 via *Agrobacterium*-mediated transformation.

**The binary vector pCGP2252 (CaMV 35S: kennedia F3'5'H: ocs 3')**

- 10 The binary vector pCGP2252 (Figure 42) contains a chimeric *CaMV 35S: kennedia F3'5'H: ocs 3'* gene in tandem with the *35S 5': SuRB* selectable marker cassette of the binary vector pCGP1988 (Figure 16).

**Intermediates in the preparation of the binary vector pCGP2252**

- 15 The plasmid pCGP2231 was firstly linearized upon digestion with the restriction endonuclease *XhoI*. The overhanging ends were repaired and then the kennedia *F3'5'H* cDNA clone was released upon digestion with the restriction endonuclease *PstI*. The ~1.7 kb fragment was ligated with the *ClaI* (repaired ends)/*PstI* ends of pCGP2105 (Figure 17). Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The resulting plasmid was  
20 designated pCGP2236.

- A 3.6 kb fragment containing the *CaMV 35S: kennedia F3'5'H: ocs 3'* chimeric gene cassette was released from the plasmid pCGP2236 upon digestion with the restriction endonucleases *XhoI* and *NotI*. The overhanging ends were repaired and the purified  
25 fragment was ligated with *Asp718* repaired ends of the binary vector, pCGP1988 (Figure 16). Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated pCGP2252 (Figure 42).

- 30 **Rose transformation with pCGP2252**

- 103 -

The T-DNA contained in the binary vector plasmid pCGP2252 (Figure 42) was introduced into *Rosa hybrida* cv. Kardinal via *Agrobacterium*-mediated transformation.

#### Butterfly pea F3'5'H constructs

##### 5 Isolation of a F3'5'H cDNA clone from petals of *Clitoria ternatea* (butterfly pea)

##### Construction of butterfly pea petal cDNA library

A blue variety of *Clitoria ternatea* (butterfly pea, the seeds were kindly provided by Osaka Botanical Garden) was grown in a field in Osaka. Total RNA was isolated from fresh and pigmented petals at a pre-anthesis stage as described above. PolyA<sup>+</sup> RNA was prepared  
10 using Oligotex (Takara) according to the manufacturer's recommendation. A petal cDNA library of butterfly pea was constructed from the polyA<sup>+</sup> RNA using a directional λZAP-cDNA synthesis kit (Stratagene, USA) following the manufacturer's protocols.

##### Screening of butterfly pea cDNA library for a F3'5'H cDNA clone

15 The butterfly pea petal cDNA library was screened with DIG-labelled petunia F3'5'H *petHf1* cDNA clone as described previously (Tanaka *et al.*, *Plant Cell Physiol.* 37: 711-716, 1996). Two cDNA clones that showed high sequence similarity to the petunia F3'5'H *petHf1* were identified. The plasmid containing the longest cDNA clone was designated pBHF2 and the cDNA clone was sequenced. Alignment between the deduced amino acid  
20 sequences of the butterfly pea F3'5'H clone and the petunia F3'5'H *petHf1* clone (SEQ ID NO:2) revealed that the butterfly pea F3'5'H cDNA (contained in pBHF2) did not represent a full-length cDNA and lacked first 2 bases of the putative initiation codon. These two bases along with a *Bam*HI restriction endonuclease recognition site were added to the cDNA clone using PCR and a synthetic primer, 5'-  
25 GGGATCCAACAATGTTCTTCTAAGAGAAAT-3' [SEQ ID NO:25] as described previously (Yonekura-Sakakibara *et al.*, *Plant Cell Physiol.* 41: 495-502, 2000). The resultant fragment was digested with the restriction endonucleases *Bam*HI and *Pst*I and the subsequent DNA fragment of about 200 bp was recovered. The DNA fragment was ligated with a 3.3 kb fragment of *Bam*HI/*Eco*RI digested pBHF2 to yield pBHF2F (Figure 43).  
30 The DNA sequence was confirmed to exclude errors made during PCR (SEQ ID NO:20).

- 104 -

Comparison of the nucleotide sequence of butterfly pea *F3'5'H* clone (SEQ ID NO:20) with that of the petunia *F3'5'H* revealed around 59% identity to the petunia *F3'5'H petHf1* clone (SEQ ID NO:1) and 62% identity to the petunia *F3'5'H petHf2* clone (SEQ ID NO:3).

5

**The binary vector pCGP2135 (*AmCHS* 5': butterfly pea *F3'5'H*: *petD8* 3')**

The plasmid pCGP2135 (Figure 44) contains the butterfly pea *F3'5'H* cDNA clone between a snapdragon *CHS* promoter fragment (*AmCHS* 5') and a petunia *PLTP* terminator fragment (*petD8* 3') in tandem with the 35S 5': *SuRB* selectable marker gene cassette of the binary vector pCGP1988 (Figure 16).

10

The petunia *F3'5'H (petHf1)* cDNA clone in pCGP725 (described in Example 4) (Figure 7) was removed by initially digesting pCGP725 with the restriction endonucleases *Xba*I and *Bam*HI. The ends were repaired the ~4.9kb fragment containing the vector with the *AmCHS* 5' and *petD8* 3' fragments was purified and ligated with the repaired ends of the ~1.6kb *Xho*I/*Bam*HI fragment from pBHF2F (Figure 43) containing the butterfly pea *F3'5'H* cDNA clone to produce pCGP2133. Correct insertion of the butterfly pea *F3'5'H* fragment in tandem with the *AmCHS* 5' and *petD8* 3' fragments was confirmed by restriction endonuclease mapping.

15

20

The *AmCHS* 5': butterfly pea *F3'5'H*: *petD8* 3' cassette was then isolated from pCGP2133 by firstly digesting with the restriction endonuclease *Not*I. The ends of the linearised plasmid were repaired and then the chimeric *F3'5'H* gene was released upon digestion with the restriction endonuclease *Eco*RV. The ~3.6kb purified fragment was then ligated with *Asp*718 repaired ends of the binary vector pCGP1988 (described in Example 4) (Figure 16). Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated pCGP2135 (Figure 44).

25



- 105 -

**Carnation and petunia transformation with pCGP2135**

The T-DNA contained in the binary vector plasmid pCGP2135 (Figure 44) was introduced into *Dianthus caryophyllus* cultivars Kortina Chanel and Monte Lisa and *Petunia hybrida* cv. Skr4 x Sw63 via *Agrobacterium*-mediated transformation.

**The binary vector pBEBF5 (eCaMV 35S: Butterfly pea F3'5'H: nos 3')**

The binary vector, pBE2113-GUS contains a *GUS* coding region between an enhanced *CaMV 35S* promoter and *nos* terminator in a pBI121 binary vector (Mitsuhara *et al.*, 1996, *supra*). The plasmid pBE2113-GUS was digested with the restriction endonuclease *SacI*. The overhanging ends were repaired and then ligated with a *SaII* linker to yield pBE2113-GUSs. The 1.8 kb *BamHI-XhoI* fragment from pBHF2F was ligated with *BamHI-SaII* digested pBE2113-GUSs to create pBEBF5 (Figure 45).

**Rose transformation with pBEBF5**

The T-DNA contained in the binary vector plasmid pBEBF5 (Figure 45) was introduced into *Rosa hybrida* cultivar Lavande via *Agrobacterium*-mediated transformation.

**The binary vector pCGP2134 (CaMV 35S: butterfly pea F3'5'H: ocs 3')**

The binary vector pCGP2134 (Figure 46) contains a chimeric *CaMV 35S: butterfly pea F3'5'H: ocs 3'* gene cassette in a tandem orientation with the *35S 5': SuRB* selectable marker gene cassette of the binary vector pCGP1988 (Figure 16).

**Intermediates in the preparation of the binary vector pCGP2134**

The butterfly pea *F3'5'H* cDNA clone was released upon digestion of the plasmid pBHF2F (Figure 43) with the restriction endonucleases *XhoI* and *BamHI*. The overhanging ends were repaired and the ~1.7 kb fragment was ligated with the *PstI* (repaired ends)/*EcoRV* ends of pCGP2105 (described in Example 4) (Figure 17). Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated pCGP2132.

- 106 -

An ~3.6 kb fragment containing the *CaMV 35S: butterfly pea F3'5'H: ocs 3'* chimeric gene cassette was released upon digestion with the restriction endonucleases *XhoI* and *XbaI*. The overhanging ends were repaired and the purified fragment was ligated with *Asp718* repaired ends of the binary vector, pCGP1988 (described in Example 4) (Figure 16).

- 5 Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated pCGP2134 (Figure 46).

**Rose transformation with pCGP2134**

- 10 The T-DNA contained in the binary vector plasmid pCGP2134 (Figure 46) was introduced into *Rosa hybrida* cv. Kardinal via *Agrobacterium*-mediated transformation.

**Gentia F3'5'H constructs**

**Isolation of a F3'5'H cDNA clone from petals of *Gentiana triflora* (gentian).**

- 15 **Construction and screening of a gentian petal cDNA library**

- The isolation of a gentian cDNA encoding *F3'5'H* has been described previously (Tanaka *et al.*, 1996, *supra*) and is contained within the plasmid pG48 (Figure 47). Comparison of the nucleotide sequence of the gentia *F3'5'H* clone (*Gen#48*) (SEQ ID NO:22) contained in the plasmid pG48 (Figure 47) with that of the petunia *F3'5'H* revealed around 61% identity to the petunia *F3'5'H petHf1* clone (SEQ ID NO:1) and 64% identity to the petunia *F3'5'H petHf2* clone (SEQ ID NO:3).
- 20

**The binary vector pCGP1498 (*AmCHS* 5': gentia *F3'5'H*: *petD8* 3')**

- The plasmid pCGP1498 (Figure 48) contains the gentia *F3'5'H* (*Gen#48*) cDNA clone between a snapdragon *CHS* promoter fragment (*AmCHS* 5') and a petunia *PLTP* terminator fragment (*petD8* 3') in tandem with the *35S* 5': *SuRB* selectable marker gene cassette of the binary vector pWTT2132 (Figure 6).
- 25

- The petunia *F3'5'H* (*petHf1*) cDNA clone in pCGP725 (described in Example 4) (Figure 7) was removed by initially digesting pCGP725 with the restriction endonucleases *XbaI* and *BamHI*. The ends were repaired the ~4.9kb fragment containing the vector with the
- 30

- 107 -

5 *AmCHS* 5' and *petD8* 3' fragments was purified and ligated with the repaired ends of the ~1.7 kb *XhoI/BamHI* fragment from pG48 (Figure 47) containing the *gentia* F3'5'H cDNA clone to produce pCGP1496. Correct insertion of the *gentia* F3'5'H fragment in tandem with the *AmCHS* 5' and *petD8* 3' fragments was confirmed by restriction endonuclease mapping.

10 The *AmCHS* 5': *gentia* F3'5'H: *petD8* 3' cassette was then isolated from pCGP1496 by firstly digesting with the restriction endonuclease *NcoI*. The overhanging ends of the linearised plasmid were repaired and then the chimeric F3'5'H gene was released upon digestion with the restriction endonuclease *EcoRV*. The ~3.6kb purified fragment was then ligated with *Asp718* repaired ends of the binary vector pWTT2132 (Figure 6). Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated pCGP1498 (Figure 48).

15

**Carnation and petunia transformation with pCGP1498**

The T-DNA contained in the binary vector plasmid pCGP1498 (Figure 48) was introduced into *Dianthus caryophyllus* cultivars Kortina Chancel and Monte Lisa and *Petunia hybrida* cv. Skr4 x Sw63 via *Agrobacterium*-mediated transformation.

20

**The binary vector pBEGHF48 (eCaMV 35S: *gentia* F3'5'H: nos 3')**

25 The *gentia* F3'5'H cDNA clone was released by digestion of the plasmid pG48 with the restriction endonucleases *BamHI* and *XhoI*. The resulting ~1.7 kb DNA fragment was isolated and ligated with *BamHI/SalI* digested pBE2113-GUSs (Mitsuhara *et al.*, 1996, *supra*) to create pBEGHF48 (Figure 49).

**Rose transformation with pBEGHF48**

The T-DNA contained in the binary vector plasmid pBEGHF48 (Figure 49) was introduced into *Rosa hybrida* cv. Lavande via *Agrobacterium*-mediated transformation.

30

- 108 -

**The binary vector pCGP1982 (CaMV 35S: gentia F3'5'H: ocs 3')**

The binary vector pCGP1982 (Figure 50) contains a chimeric *CaMV 35S: gentia F3'5'H: ocs 3'* gene cassette in tandem with the *35S 5': SuRB* selectable marker gene cassette of the binary vector pWTT2132 (Figure 6).

5

**Intermediates in the preparation of the binary vector pCGP1982**

The plasmid pG48 (Figure 47) was linearised upon digestion with the restriction endonuclease *Asp718*. The overhanging ends were repaired and then the *gentia F3'5'H* cDNA clone (*Gen#48*) was released upon digestion with the restriction endonuclease  
10 *Bam*HI. The ~1.7 kb fragment was ligated with the 5.95kb *Eco*RI (repaired ends)/*Bam*HI fragment of pKIWI101 (Janssen and Gardner, 1989, *supra*). Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated pCGP1981.

15 An ~3.6 kb fragment containing the *CaMV 35S: gentia F3'5'H: ocs 3'* chimeric gene cassette was released upon digestion of the plasmid pCGP1981 with the restriction endonucleases *Xho*I and *Xba*I. The overhanging ends were repaired and the purified fragment was ligated with repaired ends of *Asp*718 digested binary vector, pWTT2132. Correct insertion of the fragment was established by restriction endonuclease analysis of  
20 plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated pCGP1982 (Figure 50).

**Rose transformation with pCGP1982**

The T-DNA contained in the binary vector plasmid pCGP1982 (Figure 50) was introduced  
25 into *Rosa hybrida* cv. Kardinal via *Agrobacterium*-mediated transformation.

### Lavender F3'5'H constructs

#### Isolation of a F3'5'H cDNA clone from petals of *Lavandula nil* (lavender)

##### Construction of lavender petal cDNA library

Cut flowers of a violet variety of *Lavandula nil* were purchased from a florist. Total RNA  
5 was isolated from fresh and pigmented petals as described above. PolyA<sup>+</sup> RNA was  
prepared using Oligotex (Takara) according to the manufacturer's recommendations. A  
petal cDNA library of lavender was constructed from the polyA<sup>+</sup> RNA using a directional  
λZAP-cDNA synthesis kit (Stratagene, USA) following the manufacturer's protocols.

##### 10 Screening of lavender cDNA library for a F3'5'H cDNA clone

The lavender petal cDNA library was screened with DIG labelled petunia F3'5'H *petHf1*  
cDNA clone as described previously (Tanaka *et al.* 1996, *supra*). One cDNA clone (LBG)  
that showed high similarity to petunia F3'5'H *petHf1* was identified and the plasmid was  
designated pLHF8 (Figure 51). The nucleotide sequence of the lavender F3'5'H (LBG)  
15 cDNA clone was designated as SEQ ID NO: 31.

Comparison of the nucleotide sequence of lavender F3'5'H clone with that of the petunia  
F3'5'H cDNA clones revealed around 59% identity to the petunia F3'5'H *petHf1* clone  
(SEQ ID NO:1) and 60% identity to the petunia *petHf2* clone (SEQ ID NO:3).

20

##### The binary vector pBELF8 (eCaMV 35S: lavender F3'5'H; nos 3')

The plasmid of pLHF8 (Figure 51) was digested with the restriction endonucleases *Bam*HI  
and *Xho*I to release a DNA fragment of approximately 1.8 kb. The ~1.8kb purified  
fragment from pLHF8 was then ligated with the *Bam*HI-*Sa*I digested ends of the plasmid  
25 pBE2113-GUSs (described above) to create pBELF8 (Figure 52).

##### Rose transformation with pBELF8

The T-DNA contained in the binary vector plasmid pBELF8 (Figure 52) was introduced  
into *Rosa hybrida* cultivar Lavande via *Agrobacterium*-mediated transformation.

30

- 110 -

**EXAMPLE 8*****Analysis of transgenic carnation, petunia and rose***

The transgenic plants produced in the experiments described in Example 7 were grown to flowering. Flowers were collected and the colors of the petals were coded using the Royal Horticultural Society Colour Charts (RHSCC). The anthocyanins were extracted and the anthocyanidins analysed by spectrophotometric, TLC and/or HPLC analysis. Total RNA was also isolated from petal tissue of the appropriate stages of flower development and Northern blot analysis was used to detect transcripts of *F3'5'H* transgenes. The results of the transgenic analysis are summarised in Tables 11, 12 and 13.

**Carnation**

The *F3'5'H* genes described in Example 7 were evaluated for their ability to lead to the production of delphinidin-based pigments in carnation petals. Two carnation cultivars, Kortina Chanel (KC) and Monte Lisa (ML), were used in the transformation experiments.

The carnation cultivar Kortina Chanel produces pink colored flowers that normally accumulate cyanidin-based anthocyanins. This cultivar therefore contains a carnation *F3'H* and DFR activity that an introduced *F3'5'H* would need to compete with for substrate. The carnation cultivar Monte Lisa produces brick red colored flowers that normally accumulate pelargonidin-based anthocyanins. This cultivar is thought to lack fully functional *F3'H* activity and contain a DFR that is capable of acting on DHK and thus an introduced *F3'5'H* would only be required to compete with the endogenous DFR for substrate.

- 111 -

**TABLE 11** Results of transgenic analysis of petals from carnations transformed with T-DNAs containing *F3'5'H* gene expression cassettes (*AmCHS* 5': *F3'5'H*: *petD8* 3').

<i>F3'5'H</i>	pCGP	cv.	#tg	TLC+	HPLC+	Highest % del	Av. % del	Northern+
Salvia#2	2121	KC	22	2/16	3/4	12.5%	7%	nd
		ML	21	17/18	9/9	76%	57%	14/15
Salvia#47	2122	KC	23	6/12	8/8	29%	12%	nd
		ML	25	21/22	17/17	88%	56%	12/14
Sollya	2130	KC	30	22/27	17/17	35%	11%	nd
		ML	23	14/15	14/14	76%	49%	13/14
Butterfly pea	2135	KC	22	0/16	0/1	nd	nd	nd
		ML	24	19/20	13/13	23%	10%	14/14
Gentian	1498	KC	22	0/14	nd	nd	nd	7/8
		ML	2	2/2	1/1	nd	nd	1/2
pansy BP#18	1972	KC	26	18/20	12/12	14%	9%	19/19
		ML	21	15/16	8/8	80%	66%	14/16
pansy BP#40	1973	KC	26	11/15	7/8	18%	8%	13/17
		ML	33	19/22	20/20	72%	52%	12/15
petunia <i>petHf1</i>	1452	KC	104	41/64	nd	3.5%	1.3%	15/17
		ML	48	39/41	26/26	75%	30%	12/13
petunia <i>petHf2</i>	1524	ML	27	18/19	17/17	81%	41%	12/14

- 5 *F3'5'H* = *F3'5'H* sequence contained on the T-DNA
- pCGP = plasmid pCGP number of the binary vector used in the transformation experiment
- cv. = cultivar
- 10 KC = carnation cultivar Kortina Chanel (cyanidin line)
- ML = carnation cultivar Monte Lisa (pelargonidin line)
- #tg = total number of transgenics produced

- 112 -

TLC+ = number of individual events in which delphinidin or delphinidin-based molecules was detected in petals (as determined by TLC) over the total number of individual events analyzed

HPLC+ = number of individual events in which delphinidin or delphinidin-based molecules was detected in petals (as determined by HPLC) over the total number of individual events analyzed

Highest % del = Highest % delphinidin or delphinidin-based molecules detected in the petals for the population of transgenic events

Av % del = average % delphinidin or delphinidin-based molecules detected in the petals for the population of transgenic events

Northern = number of individual events in which the specific intact *F3'5'H* transcripts were detected by Northern blot analysis in total RNA isolated from petals over the total number of events analyzed

*nd* = not done

The results suggest that all of the *F3'5'H* sequences evaluated (petunia *petHf1*, petunia *petHf2*, *Salvia Sal#2*, *Salvia Sal#47*, *Solysa Sol#5*, Butterfly pea *BpeaHF2*, pansy *BP#18*, pansy *BP#40* and *Gentian Gen#48*) were stable in carnation and resulted in the production of novel delphinidin-based pigments in carnation flowers. Intact transcripts of each *F3'5'H* were detected by Northern blot analysis in total RNA isolated from petals of the transgenic carnations.

#### Petunia

The *F3'5'H* genes described in Example 7 were evaluated for their ability to lead to the production of delphinidin-based pigments in petunia petals. The *P. hybrida* F1 hybrid Skr4 x SW63 which is homozygous recessive for *Hf1* and *Hf2*, was used in the transformation experiments. Although Skr4 x SW63 is homozygous recessive for *Hf1* and *Hf2*, these mutations do not completely block production of the endogenous *F3'5'H* (see US Patent Number 5,349,125) and low levels of malvidin are produced to give the petal limb a pale lilac color. Malvidin is the methylated derivative of the 3'5'-hydroxylated pigment, delphinidin or delphinidin-based molecules (Figures 1A and 1B). Spectrophotometric



- 113 -

analysis was used as a measure of total anthocyanins accumulating in petals from the transgenic petunia flowers. The increased level of anthocyanins and/or the color change detected was used as a guide to the efficacy of the *F3'5'H* gene under evaluation.

5 **TABLE 12** Results of transgenic analysis of petals from *P. hybrida* cv Skr4 x SW63 plants transformed with T-DNAs containing *F3'5'H* gene expression cassettes (*AmCHS* 5': *F3'5'H*: *petD8* 3').

<b>F3'5'H</b>	<b>pCGP</b>	<b># tg</b>	<b>TLC+</b>	<b>Col</b>	<b>† A/c</b>	<b>Best</b>	<b>Av.</b>	<b>Northern+</b>	<b>Best color</b>
control	na	na	na	na	na	144-250		0	75C
Gentian#48	1498	22	3/5	18/20	nd			6/6	72B/78A
Butterfly pea	2135	24	18/20	22/24	23/24	4427	2397	nd	74A/78A
Kennedia	2256	24	22/24	22/24	22/24	4212	2592	nd	74A/78A
Salvia#2	2121	24	21/24	21/24	21/24	2471	1730	nd	78A
Salvia#47	2122	19	17/19	16/19	16/19	2634	1755	nd	78A/80A
Sollya#5	2130	22	14/16	13/16	13/16	3446	1565	nd	78A
pansy BP#18	1972	22	nd	20/22	nd	nd	nd	9/9	74A/B
pansy BP#40	1973	19	8/8	18/19	18/20	2583	1556	nd	74/78A
petunia <i>petHf1</i>	484	16	nd	9/16	8/15	2683	1250	nd	74A/B
petunia <i>petHf2</i>	1524	20	nd	18/20	8/8	4578	2357	8/8	74A/B

- 10 **F3'5'H** = *F3'5'H* sequence contained on the T-DNA  
**pCGP** = plasmid pCGP number of the binary vector used in the transformation experiment  
**#tg** = total number of transgenics produced

- 114 -

TLC+ = number of individual events in which malvidin was detected in the flowers (at a level above the Skr4 x Sw63 background) (as determined by TLC) over the total number of individual events analyzed

Col = number of individual events that produced flowers with an altered flower color compared to the control over the total number examined

† A/c = number of individual events that had an increased level of anthocyanins in petals as measured by spectrophotometric analysis of crude extracts over the number of individual events analyzed (in  $\mu\text{moles/g}$ )

Best = highest anthocyanin amount as measured by spectrophotometric analysis of crude extracts from a flower of an individual event (in  $\mu\text{moles/g}$ )

Av = the average amount of anthocyanin detected as measured by spectrophotometric analysis of crude extracts from a flower in the population of transgenic flowers analysed (in  $\mu\text{moles/g}$ )

Northern = number of individual events in which the specific intact *F3'5'H* transcripts were detected by Northern blot analysis in total RNA isolated from petals over the total number of events analyzed

Best color = most dramatic color change recorded for the transgenic population

nd = not done

na = not applicable

Introduction of the *F3'5'H* gene expression cassettes into Skr4 x SW63 led to a dramatic flower color change from pale lilac to purple with a dramatic increase in the production of malvidin in the petals..

The results suggest that all of the *F3'5'H* sequences tested (petunia *petHf1*, petunia *petHf2*, *Salvia Sal#2*, *Salvia Sal#47*, *Sollya Sol#5*, Butterfly pea *BpeaHF2*, pansy *BP#18*, pansy *BP#40*, *Gentian Gen#48*, *Kennedia Kenn#31*) were stable in petunia petals and resulted in the complementation of the *Hf1* or *Hf2* mutation in the Skr4 x SW63 petunia line leading to dramatically increased levels of malvidin accumulation with a concomitant color change

## Rose

The *F3'5'H* genes described in Example 7 were evaluated for their ability to lead to the production of delphinidin-based pigments in rose petals. A selection of three rose cultivars, Kardinal (Kard), Soft Promise (SP) or Lavande (Lav) were used in transformation experiments. The rose cultivar Kardinal produces red colored flowers that normally accumulate cyanidin-based anthocyanins. This cultivar therefore contains rose *F3'H* and DFR activities that the introduced *F3'5'H* would need to compete with for substrate. The rose cultivar Lavande produces light pink colored flowers that normally accumulate cyanidin-based anthocyanins. This cultivar therefore contains functional rose *F3'H* and DFR activities that the introduced *F3'5'H* would need to compete with for substrate. The rose cultivar Soft Promise produces apricot colored flowers that normally accumulate pelargonidin. This cultivar is thought to lack a fully functional rose *F3'H* activity and contain a DFR that is capable of acting on DHK and thus the introduced *F3'5'H* would only be required to compete with the endogenous rose DFR for substrate.

**TABLE 13** Results of transgenic analysis of petals from roses transformed with T-DNAs containing *F3'5'H* gene expression cassettes (*CaMV 35S: F3'5'H: ocs 3'*).

<i>F3'5'H</i>	plasmid	Cult	#tg	TLC +	HPLC+	Highest % del	Av. % del	Northern+
Salvia2	pCGP2120	Kard	30	18/20	21/21	12%	5%	18/18
Salvia47	pCGP2119	Kard	22	11/16	9/9	7.1%	2%	12/15
Sollya	pCGP2131	Kard	27	0/23	2/2	1%	0.5%	6/6
Butterfly pea	pCGP2134	Kard	29	0/15	nd	na	na	0/9
	pBEBF5	Lav	25	nd	0/25	0%	0%	nd
Gentian	pCGP1482	Kard	27	0/23	nd	na	na	0/23
	pBEGHF48	Lav	23	nd	0/23	0%	0%	0/23
pansy BP18	pCGP1967	Kard	56	30/33	33/34	58%	12%	21/21
		SP	36	21/24	18/18	65%	35%	16/21
pansy BP40	pCGP1969	Kard	22	15/15	15/15	24%	9%	16/16
		SP	37	17/17	16/17	80%	54%	11/13

- 116 -

<i>F3'5'H</i>	plasmid	Cult	#tg	TLC +	HPLC+	Highest % del	Av. % del	Northern+
Petunia	pCGP1638	Kard	22	0/21	nd	na	na	0/16
<i>petHf1</i>	pCGP1392	Lav	34	nd	0/34	0%	0%	nd
Petunia	pCGP2123	Kard	41	0/26	nd	na	na	0/10
<i>petHf2</i>								
Lavender	pBELF8	Lav	28	nd	4/28	4%	3.5%	nd

*F3'5'H* = the *F3'5'H* sequence contained on the T-DNA

plasmid = the plasmid number of the binary vector used in the transformation experiment

5 Cult = *Rosa hybrida* cultivar

Kard = Kardinal

SP = Soft Promise

Lav = Lavande

#tg = # of independent transgenic events produced

10 TLC+ = number of individual events that accumulated detectable delphinidin or delphinidin-based molecules (as determined by TLC) in the petals over the number of individual events analyzed

HPLC+ = number of individual events that accumulated detectable delphinidin or delphinidin-based molecules (as determined by HPLC) in the petals over the number of individual events analyzed

15 Northern = number of individual events in which the specific intact *F3'5'H* transcripts were detected by Northern blot analysis in total RNA isolated from petals over the total number of events analyzed

20 nd = not done

- 117 -

The results suggest surprisingly that not all of the *F3'5'H* sequences assessed (petunia *petHf1*, petunia *petHf2*, Salvia *Sal#2*, Salvia *Sal#47*, Sollya *Sol#5*, Butterfly pea *BpeaHF2*, pansy *BP#18*, pansy *BP#40*, Gentian *Gen#48*, Kennedia *Kenn#31* and Lavender *LBG*) were functional in rose. In fact transcripts of the introduced *F3'5'H* sequences isolated

5 from *Clitoria ternatea* (butterfly pea), *Gentiana triflora*, (gentian) and *Petunia hybrida* (petunia) failed to accumulate in rose petals. Only full-length *F3'5'H* transcripts from pansy, salvia, kennedia, sollya and lavender accumulated in rose petals. However although Kennedia *F3'5'H* transcripts did accumulate in rose petals, there was either no

10 accumulation of the enzyme or the enzyme produced was either not functional or was unable to compete with the endogenous rose *F3'H* and *DFR* enzymes to allow for the production of delphinidin or delphinidin-based molecules pigments. Of the *F3'5'H* sequences evaluated, only the *F3'5'H* sequences derived from cDNA clones from *Salvia* spp. (*Sal#2* and *Sal#47*), *Viola* spp. (*BP#18* and *BP#40*), *Sollya* spp. (*Sol#5*) and *Lavandula nil* (*LBG*) resulted in the production of delphinidin or delphinidin-based

15 molecules based pigments in rose petals. Based on the relative percentages of delphinidin or delphinidin-based molecules produced in rose petals, the *F3'5'H* sequences from pansy (*BP#18* and *BP#40*) were revealed to be the most effective of those assessed at producing delphinidin or delphinidin-based molecules in rose petals.

20 **Introduction of *Viola* spp. *F3'5'H* sequence into *Rosa hybrida* cv. Medeo and Pamela**

As described in the introduction, copigmentation with other flavonoids, further modification of the anthocyanidin molecule and the pH of the vacuole impact on the color produced by anthocyanins. Therefore, selection of rose cultivars with relatively high levels of flavonols and relatively high vacuolar pH would result in bluer flower colors upon

25 production of delphinidin or delphinidin-based molecules pigments.

The rose cultivar Medeo generally produces cream-colored to pale apricot flowers (RHSCC 158C to 159A). HPLC analysis of the anthocyanidins and flavonols accumulating in Medeo rose petals revealed that the petals accumulate high levels of flavonols (2.32

30 mg/g kaempferol, 0.03 mg/g quercetin) and very low levels of anthocyanins (0.004 mg/g

- 118 -

cyanidin, 0.004 mg/g pelargonidin). The estimated vacuolar pH of Medeo petals is around 4.6.

5 The rose cultivar Pamela produces white to very pale pink colored flowers. It similarly accumulates low levels of anthocyanin and relatively high levels of flavonols.

The T-DNA contained in the construct pCGP1969 (Figure 30) incorporating the pansy *F3'5'H* clone, *BP#40*, was also introduced into the rose cultivars Medeo and Pamela resulting in the production of over 90% delphinidin or delphinidin-based molecules in  
10 these roses and leading to a dramatic color change and novel colored flowers. The most dramatic color change in transgenic Medeo flowers was to a purple/violet color of RHSCC 70b, 70c, 80c, 186b. The most dramatic color change in transgenic Pamela flowers was to a purple/violet color of RHSCC 71c, 60c, 71a, 80b.

15 In conclusion, two unexpected findings were revealed when gene sequences that had been proven to lead to functionality in petunia and carnation were introduced into roses.

First, the petunia *F3'5'H* *petHf1* (and *petHf2*) sequences that had resulted in novel color production in carnation and also proven to lead to synthesis of a functional enzyme in  
20 petunia did not lead to full-length (or intact) transcript accumulation (as detectable by Northern blot analysis) in rose petals. In fact, there was either no accumulation of full-length or intact transcript or the transcripts that were detected were degraded and were seen as low MW (or fast migrating) smears on RNA blots indicating the presence of low MW heterologous hybridizing RNA. Therefore in order to find a *F3'5'H* sequence that  
25 would accumulate in rose and lead to a functional enzyme, a number of *F3'5'H* sequences were isolated. Again it was not obvious which sequence would lead to an active enzyme in rose petals. All of the *F3'5'H* sequences isolated were tested for functionality in carnation and/or petunia and all led to accumulation of intact transcripts and production of a functional *F3'5'H* activity. However only *F3'5'H* sequences from pansy (*BP#18* and  
30 *BP#40*), salvia (*Sal#2* and *Sal#47*), sollya (*Soll#5*), kennedia (*Kenn#31*) and lavender (*LBG*) resulted in accumulation of intact full-length transcripts and only those from pansy

- 119 -

(BP#18 and BP#40), salvia (Sal#2 and Sal#47), sollya (Sol#5) and lavender (LBG) resulted in production of a functional enzyme in rose as measured by the synthesis of delphinidin or delphinidin-based molecules.

- 5 Secondly that it was not obvious which promoters would be effective in rose. Promoter cassettes that had been tested and proven to be functional in carnation and petunia flowers did not lead to accumulation of detectable transcripts in rose petals. Of the promoters tested in rose, only *CaMV 35S*, *RoseCHS 5'*, *ChrysCHS 5'*, *mas 5'* and *nos 5'* promoters led to intact and detectable *GUS* or *nptII* or *SuRB* transcript accumulation in rose petals.

10

Table 14 shows a summary of the results obtained when assessing F3'5'H sequences from various species in petunia, carnation and rose.

15 **TABLE 14** Summary of effectiveness of the F3'5'H sequences in petunia, carnation and rose

F3'5'H	Petunia		Carnation		Rose	
	Mal	RNA	Del	RNA	Del	RNA
Kennedia ( <i>Kenn#31</i> )	+	nd	nd	nd	-	+
Gentian ( <i>Gen#48</i> )	+	+	+	+	-	-
Salvia ( <i>Sal#2</i> )	+	nd	+	+	+	+
Salvia ( <i>Sal#47</i> )	+	nd	+	+	+	+
Sollya ( <i>Sol#5</i> )	+	nd	+	+	+	+
Butterfly pea ( <i>BpeaHF2</i> )	+	nd	+	+	-	-
Pansy ( <i>BP#18</i> )	+	+	+	+	+	+
Pansy ( <i>BP#40</i> )	+	nd	+	+	+	+
Petunia ( <i>petHf1</i> )	+	+	+	+	-	-
Petunia ( <i>petHf2</i> )	+	+	+	+	-	-
Lavender ( <i>LBG</i> )	nd	nd	nd	nd	nd	+

- 120 -

*nd* = not done

Mal = malvidin detected in petals as analysed by TLC

Del = delphinidin or delphinidin-based molecules detected in petals as analysed by TLC or HPLC

5 + = yes

- = no

### EXAMPLE 9

#### *Use of pansy F3'5'H sequences in species other than rose*

#### 10 *Gerbera*

From the examples above, it was clear that the pansy *F3'5'H* sequences, *BP#18* and *BP#40*, resulted in functional *F3'5'H* activity and lead to the production of high levels of delphinidin or delphinidin-based molecules in roses and carnations.

15 The T-DNA from binary construct pCGP1969 (described in Example 8) (Figure 30) containing the chimeric *CaMV 35S: pansy BP#40 F3'5'H: ocs 3'* gene expression cassette was introduced into the gerbera cultivar Boogie via *Agrobacterium*-mediated transformation, to test the functionality of the pansy *F3'5'H* sequence in gerbera.

20 Of six events produced to date, one (#23407) has produced flowers with a dramatic color change (RHSCC 70c) compared to the control flower color (RHSCC 38a, 38c).

The color change of the petals of the transgenic gerbera has been correlated with the presence of delphinidin or delphinidin-based molecules as detected by TLC.

25

#### *Other species*

In order to produce delphinidin or delphinidin-based molecules pigments in plants that do not normally produce delphinidin-based pigments and does not contain a flavonoid 3'5'-hydroxylase constructs containing a *F3'5'H* gene (such as but not limited to a chimaeric

30 *Viola spp.* and/or *Salvia spp.* and/or *Solhya spp.* and/or *Lavandula spp.* and/or *Kennedia spp.* *F3'5'H* gene) are introduced into a species that does not normally produce



- 121 -

delphinidin-based pigments. Such plants may include but are not limited to carnation, chrysanthemum, gerbera, orchids, *Euphorbia*, *Begonia* and *apple*.

#### EXAMPLE 10

5

##### Characteristics of F3'5'H sequences evaluated in *petunia*, *carnation* and *rose*

Gene regulation in eukaryotes is, in simple terms, facilitated by a number of factors which interact with a range of sequences proximal and distal to a nucleotide sequence coding for a given polypeptide. Engineering expression cassettes for introduction into plants for the generation of one or more traits is based on an understanding of gene regulation in eukaryotes in general and, in selected cases, plants in particular. The essential elements include a series of transcriptional regulation sequences typically, but not exclusively, located upstream or 5' to the point of transcription initiation. Such elements are typically described as enhancers and promoters, the latter being proximal to the point of transcription initiation. Immediately downstream from, or 3' to, the initiation of transcription point is a variable region of transcribed DNA which is denoted as the 5' untranslated region (5'utr) which plays a role in transcript stability and translational efficiency. Such sequences, when engineered into expression cassettes, are frequently chimeric and may be derived from sequences naturally occurring adjacent to the coding sequence and/or adjacent to a given promoter sequence. The coding sequence (sometimes disrupted by introns) lies 3' to the 5'utr followed by a 3'utr important to transcript (mRNA) stability and translational efficiency. Sequences 3' to the end of the coding region and 3' to the 3'utr itself are denoted as terminator sequences. All these elements make up an expression cassette. In making direct comparisons between promoters or other elements it is important to maintain uniformity in the remaining elements of an expression cassette. Hence, when comparing the efficacy of various F3'5'H sequences it was possible to confine the sequences leading to instability and the subsequent autodegradation of engineered mRNA and resultant absence of tri-hydroxylated products (delphinidin or delphinidin-based molecules derivatives) to the region coding for the F3'5'H and not to other elements in the expression cassette such as 5' utr and/or 3'utr sequences for example.

- 122 -

In an attempt to identify motifs or similarities between the *F3'5'H* sequences that resulted in full-length transcripts being detected in total RNA isolated from rose flowers, and ultimately delphinidin or delphinidin-based molecules production, comparisons across a range of parameters were performed. These included sequence identities at nucleic acid and amino acid levels, sequence alignments, taxonomic classifications, % of A or T nucleotides present in the sequence, % of codons with an A or T in the third position etc.

#### *Taxonomic classification*

- 10 The taxonomy of each species from which the *F3'5'H* sequences were isolated was examined (Table 15). There appeared to be no obvious link between the subclass classification and whether the *F3'5'H* sequence resulted in an intact transcript and subsequent delphinidin or delphinidin-based molecules production in roses.
- 15 **Table 15:** Taxonomic classifications of the species that *F3'5'H* sequences were isolated from and whether the use of the sequences resulted in intact transcript in rose petals that were detectable by RNA blot analysis.

Flower	Species	Family	Order	Subclass	Intact transcript	Delphinidin in rose petals
gentian	<i>Gentiana triflora</i>	Gentianaceae	Gentianales	Asteridae	NO	NO
lavender	<i>Lavandula nil</i>	Lamiaceae	Lamiales	Asteridae	YES	YES
salvia	<i>Salvia spp.</i>	Lamiaceae	Lamiales	Asteridae	YES	YES
sollya	<i>Sollya spp.</i>	Pittosporaceae	Apiales	Asteridae	YES	YES
petunia	<i>Petunia hybrida</i>	Solanaceae	Solanales	Asteridae	NO	NO
kennedia	<i>Kennedia spp.</i>	Fabaceae	Fabales	Rosidae	YES	NO
butterfly pea	<i>Clitoria ternatea</i>	Fabaceae	Fabales	Rosidae	NO	NO
pansy	<i>Viola spp.</i>	Violaceae	Malpighiales	Rosidae	YES	YES
rose	<i>Rosa hybrida</i>	Rosaceae	Rosales	Rosidae	na	na

**Intact transcript** = **full-length F3'5'H mRNA detected by Northern blot analysis**  
**in total RNA isolated from petals from transgenic roses**

### Comparison of F3'5'H nucleotide sequences

5 The nucleotide sequence identities between each of the *F3'5'H* sequences evaluated were determined using the ClustalW program (Thompson *et al.*, 1994, *supra*) within the MacVector™ version 6.5.3 application program (Oxford Molecular Ltd., England) (Table 16). There were no obvious differences between the *F3'5'H* sequences that resulted in the detection of intact full-length transcripts in RNA isolated from rose petals and those that  
10 didn't.

**Table 16: Percentage of nucleic acid sequence identity between the nucleotide sequences of the *F3'5'H* isolated from various species. *F3'5'H* sequences that resulted in intact transcripts being detected in RNA isolated from rose petals are underlined and in italics.**

[illegible]

- 124 -

***Comparison of F3'5'H translated nucleotide sequences***

The translated nucleotide sequence identities and similarities between each of the F3'5'H sequences evaluated were also determined using the ClustalW program (Thompson *et al.*, 1994, *supra*) within the MacVector™ version 6.5.3 application program (Oxford Molecular Ltd., England) (Table 17). There were no obvious differences between the F3'5'H sequences that resulted in the detection of intact full-length transcripts in RNA isolated from rose petals and those that didn't.

Table 17: Percentage of the amino acid sequence identity and similarity (in brackets) between F3'5'H sequences isolated from various species. F3'5'H sequences that resulted in intact transcripts being detected in RNA isolated from rose petals are underlined and in italics.

	<u><i>BP18</i></u>	<u><i>BP40</i></u>	<u><i>Lav</i></u>	<u><i>Sal47</i></u>	<u><i>Sal2</i></u>	<u><i>Soll</i></u>	<u><i>Kenn</i></u>	Bpea	Gent	PetHf1	PetHf2
<u><i>BP18</i></u>	100	91 (94)	65 (77)	65 (76)	65 (76)	44 (63)	69 (83)	64 (75)	69 (80)	74 (85)	74 (85)
<u><i>BP40</i></u>		100	67 (89)	66 (77)	66 (77)	46 (64)	69 (82)	64 (75)	68 (79)	74 (85)	75 (86)
<u><i>Lav</i></u>			100	75 (86)	75 (86)	45 (63)	63 (79)	59 (74)	66 (80)	68 (82)	69 (83)
<u><i>Sal47</i></u>				100	98	45 (65)	64 (78)	60 (72)	64 (76)	68 (79)	69 (81)
<u><i>Sal2</i></u>					100	45 (65)	64 (78)	60 (72)	63 (75)	68 (79)	69 (81)
<u><i>Soll</i></u>						100	46 (66)	41 (61)	44 (62)	46 (67)	46 (66)
<u><i>Kenn</i></u>							100	72 (80)	65 (75)	71 (83)	72 (83)
Bpea								100	69 (81)	65 (75)	65 (74)
Gent									100	73 (82)	73 (82)
PetHf1										100	93 (95)
PetHf2											100

- 125 -

*Percentage of nucleotides A or T in the F3'5'H DNA sequences*

There is some evidence to suggest that the choice of codons influences the rate of translation and mRNA degradation. Certain codons are used less frequently than others are and this may be related to the abundance of isoaccepting tRNAs. Transfer RNAs corresponding to rare codons are less abundant in *E.coli* and yeast than tRNAs corresponding to preferred codons (van Hoof and Green, *Plant Molecular Biology*, 35: 383-387, 1997). Examples of altering codon usage and making a gene more "plant-like" are the bacterial *B.t.* toxin gene (reviewed in Diehn *et al.*, *Genet Engin.*, 18: 83-99, 1996) and the jellyfish *gfp* gene (Haseloff *et al.*, *Proc. Natl. Acad. Sci USA*, 94: 2122-2127, 1997). However as commented in van Hoof and Green, (1997) (*supra*), the effect of eliminating the rare codons in the *B.t.* genes increased the GC content, thereby eliminating AU-rich sequences that may be responsible for improper recognition of introns and polyadenylation sites as well as removing instability determinants. Alteration of codon usage in the jellyfish *gfp* gene also resulted in removal of a cryptic intron (Haseloff *et al.*, 1996, *supra*). Studies examining the effect of codon usage and instability elements have generally been limited to differences between genes isolated from species in different kingdoms ie. bacterial versus yeast versus animal versus plant. Within the plant kingdom, differences have been observed between the dicotyledons and the monocotyledons. Studies on transgenic plants have suggested that promoter fragments used to drive gene expression in dicotyledonous plants are not as effective when used in monocotyledonous plants (see Galun and Breiman, *Transgenic Plants*, Imperial College Press, London, England, 1997). Differences in the methylation and ultimate expression of a *DFR* transgene in *Petunia hybrida* (dicot) were detected when a maize (monocot) *DFR* cDNA was compared with a gerbera (dicot) *DFR* cDNA (Elomaa *et al.*, *Molecular and General Genetics*, 248: 649-656, 1995). The conclusion was that the gerbera *DFR* cDNA had a higher AT content (lower GC content) and was more "compatible" with the genomic organization of petunia preventing it being recognised as a foreign gene and hence silenced by methylation. (Rose along with carnation and petunia are dicotyledons and the *F3'5'H* genes tested were all isolated from dicotyledonous plants.) These points serve to illustrate that degradation and stability mechanisms are not understood in detail and differences appear between plants and other kingdoms and within the plant kingdom.

- 126 -

The content of A and T was examined in the *F3'5'H* cDNAs evaluated along with that of four flavonoid pathway genes (*F3'H*, *DFR*, *CHS*, *FLS*) that had been isolated from rose (Table 18). The third position of each codon (within the open reading frame) was also  
 5 examined and the percentage of codons with an A or a T in the third position was calculated (Table 18).

Table 18: Summary of the percentage amount of A or T dinucleotides in the *F3'5'H* sequences isolated and whether the *F3'5'H* resulted in full-length transcripts being  
 10 detected in rose petals by Northern blot analysis.

<b>F3'5'H seq</b>	<b>%AT</b>	<b>% A or T in 3rd</b>	<b>RNA</b>	<b>Delphinidin</b>
Viola BP#18	50	40	YES	YES
Viola BP#40	51	35	YES	YES
Salvia#2	48	33	YES	YES
Salvia#47	48	34	YES	YES
Sollya#5	54	54	YES	YES
LavenderLBG	50	37	YES*	YES
Kennedia#31	54	47	YES	NO
petunia <i>petHf1</i>	61	66	NO	NO
petunia <i>petHf2</i>	59	65	NO	NO
Gentian#48	57	57	NO	NO
Butterfly pea# <i>HF2</i>	57	53	NO	NO
rose <i>F3'H</i>	47	34	**	na
rose <i>CHS</i>	52	42	**	na

- 127 -

F3'5'H seq	%AT	% A or T in 3rd	RNA	Delphinidin
rose <i>DFR</i>	53	46	**	na
rose <i>FLS</i>	56	43	**	na

%AT = % of nucleotides that are A or T in the nucleic acid sequence

%A or T in 3<sup>rd</sup> = the percentage of codons that have an A or T in the third position

RNA = whether a full-length mRNA transcript was detected by Northern blot analysis in total RNA isolated from rose petals

Del = whether any *delphinidin* or *delphinidin-based molecules* was detected by TLC or HPLC in rose petals

YES\*

=although Northern blot analysis of transgenic

roses transformed with the lavender F3'5'H expression cassettes was not performed, it can be assumed that full-length transcript was produced since *delphinidin* or *delphinidin-based molecules* was detected in the rose petals.

rose F3'H (described in International Patent Application No. PCT/AU97/00124)

rose *DFR* (Tanaka et al., 1995, *supra*)

rose *FLS* (GenBank accession number AB038247)

15 rose *CHS* (GenBank accession number AB038246)

The AT content of the four rose sequences (above) encoding flavonoid pathway enzymes had an AT content of between 47 and 56%. In general the AT content of the F3'5'H sequences that resulted in intact transcripts in rose petals was between 48 and 54%.

20 However the F3'5'H sequences that did not result in intact transcripts accumulating in rose petals generally had a higher AT content of between 57 and 61%. Hence the AT content of the introduced F3'5'H genes into rose may be a factor in whether an intact transcript accumulates in rose petals and so leads to production of F3'5'H and *delphinidin* or *delphinidin-based molecules*.

25

- 128 -

The nucleotide base at the third position of each codon of the four rose sequences encoding flavonoid pathway enzymes was generally an A or a T in 34 to 46% of the codons. In general *F3'5'H* sequences that resulted in intact transcripts in rose petals contained an A or a T in the third position of each codon in 33 to 54% of the codons. However the *F3'5'H* sequences that did not result in intact transcripts accumulating in rose petals generally contained an A or a T in the third position of each codon in 53 to 66% of the codons. So the percentage of codons with an A or a T in the third position of the introduced *F3'5'H* genes into rose may also be a factor in whether an intact transcript is accumulates in rose petals and so leads to production of *F3'5'H* and *delphinidin* or *delphinidin-based* molecules.

It may be that by altering the overall content of the nucleotides A and/or T in any *F3'5'H* DNA sequence that does not result in an intact transcript in rose such as but not limited to the *Petunia hybrida petHf1*, *Petunia hybrida petHf2*, *Clitoria ternatea* (butterfly pea) *BpeaHF2* or *Gentiana triflora* (gentian) *Gen#48*, to a level more consistent with that found in rose genes, intact transcripts will accumulate and result in the efficient translation of *F3'5'H* transcripts and so to *delphinidin* or *delphinidin-based* molecules accumulation in rose petals. One way of altering the AT content of the DNA sequence without altering the amino acid sequence is to target the degeneracy of the third position of each codon.

Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the invention includes all such variations and modifications. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all combinations of any two or more of said steps or features.



**BIBLIOGRAPHY**

*A look beyond transcription*, ASPP, USA, Bailey-Serres and Gallie, eds, 1998

Altschul *et al.*, *J. Mol. Biol.* 215(3): 403-410, 1990.

Altschul *et al.*, *Nucl. Acids Res.* 25: 3389-3402, 1997.

Ausubel *et al.*, "Current Protocols in Molecular Biology" John Wiley & Sons Inc, 1994-1998, Chapter 15.

Aviv and Leder, *Proc. Natl. Acad. Sci. USA* 69: 1408, 1972.

Bevan, *Nucleic Acids Res* 12: 8711-8721, 1984.

Bonner and Laskey, *Eur. J. Biochem.* 46: 83, 1974.

Bodeau, Molecular and genetic regulation of Bronze-2 and other maize anthocyanin genes. Dissertation, Stanford University, USA, 1994

Brendel *et al.*, In *A look beyond transcription*, ASPP, USA, Bailey-Serres and Gallie, eds, 1998.

Brouillard and Dangles, In: *The Flavonoids -Advances in Research since 1986*. Harborne, J.B. (ed), Chapman and Hall, London, UK, 1-22, 1993.

Brugliera *et al.*, *Plant J.* 5: 81-92, 1994.

Brugliera, Characterization of floral specific genes isolated from *Petunia hybrida*. RMIT, Australia. PhD thesis, 1994.

- 130 -

Bullock *et al.*, *Biotechniques* 5: 376, 1987.

Comai *et al.*, *Plant Mol. Biol.* 15: 373-381, 1990.

Depicker *et al.*, *J Mol. and Appl. Genetics* 1: 561-573, 1982.

Diehn *et al.*, *Genet Engin.* 18: 83-99, 1996

Elomaa *et al.*, *Molecular and General Genetics*, 248: 649-656, 1995

Franck *et al.*, *Cell* 21: 285-294, 1980.

Galun and Breiman, *Transgenic Plants*, Imperial College Press, London, England, 1997

Guilley *et al.*, *Cell* 30: 763-773, 1982.

Hanahan, *J. Mol. Biol.* 166: 557, 1983.

Harpster *et al.*, *MGG* 212: 182-190, 1988.

Haseloff *et al.*, *Proc. Natl. Acad. Sci USA*, 94: 2122-2127, 1997

Holton and Cornish, *Plant Cell* 7: 1071-1083, 1995.

Holton and Graham, *Nuc. Acids Res.* 19: 1156, 1990.

Holton *et al.*, *Nature* 366: 276-279, 1993a.

Holton *et al.*, *Plant J.* 4: 1003-1010, 1993b

Holton, Isolation and characterization of petal specific genes from *Petunia hybrida*. PhD thesis, University of Melbourne, Australia, 1992

Huang and Miller, *Adv. Appl. Math.* 12: 373-381, 1991.

Inoue *et al.*, *Gene* 96: 23-28, 1990.

Janssen and Gardner, *Plant Molecular Biology*, 14: 61-72, 1989

Jefferson, *et al.*, *EMBO J.* 6: 3901-3907, 1987.

Johnson *et al.*, *In A look beyond transcription*, ASPP, USA, Bailey-Serres and Gallie, eds, 1998.

Karn *et al.*, *Gene* 32: 217-224, 1984.

Lazo *et al.* *Bio/Technology* 9: 963-967, 1991.

Lee *et al.*, *EMBO J.* 7: 1241-1248, 1988.

Lu *et al.*, *Bio/Technology* 9: 864-868, 1991.

Marchant *et al.*, *Molecular Breeding* 4: 187-194, 1998.

Marmur and Doty, *J. Mol. Biol.* 5: 109, 1962.

McBride and Summerfelt, *Plant Molecular Biology* 14: 269-276, 1990.

Merrifield, *J. Am. Chem. Soc.* 85: 2149, 1964.

Mitsuhara *et al.*, *Plant Cell Physiol.* 37: 49-59, 1996.

- 132 -

Mol *et al.*, *Trends Plant Sci.* 3: 212-217, 1998.

Pearson and Lipman, *Proc. Natl. Acad. Sci. USA* 85(8): 2444-2448, 1988.

Plant Molecular Biology Labfax, Croy (ed), Bios scientific Publishers, Oxford, UK, 1993.

Plant Molecular Biology Manual (2<sup>nd</sup> edition), Gelvin and Schilperoot (eds), Kluwer Academic Publisher, The Netherlands, 1994.

Robinson and Firoozabady, *Scientia Horticulturae* 55: 83-99, 1993.

Rout *et al.*, *Scientia Horticulturae* 81: 201-238, 1999.

Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, USA, 1989.

Sambrook and Russell, *Molecular Cloning: A Laboratory Manual* 3<sup>rd</sup> edition, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, USA, 2001

Seitz and Hinderer, Anthocyanins. In: *Cell Culture and Somatic Cell Genetics of Plants*. Constabel, F. and Vasil, I.K. (eds.), Academic Press, New York, USA, 5: 49-76, 1988.

Short *et al.*, *Nucl. Acids Res.* 16: 7583-7600, 1988.

Sommer and Saedler, *Mol Gen. Gent.*, 202: 429-434, 1986.

Strack and Wray, In: *The Flavonoids -Advances in Research since 1986*. Harborne, J.B. (ed), Chapman and Hall, London, UK, 1-22, 1993.

Tanaka *et al.*, *Plant Cell Physiol.* 36: 1023-1031, 1995.

- 133 -

Tanaka *et al.*, *Plant Cell Physiol.* 37: 711-716, 1996.

Thompson *et al.*, *Nucleic Acids Research* 22: 4673-4680, 1994.

Turpen and Griffith, *BioTechniques* 4: 11-15, 1986.

van Hoof and Green, *Plant Molecular Biology*, 35: 383-387, 1997

Winkel-Shirley, *Plant Physiol.* 126: 485-493, 2001a.

Winkel-Shirley, *Plant Physiol.* 127: 1399-1404, 2001b.

Yonekura-Sakakibara *et al.*, *Plant Cell Physiol.* 41: 495-502, 2000.